

RECIPROCITY SHEAVES, II

BRUNO KAHN, SHUJI SAITO, AND TAKAO YAMAZAKI

ABSTRACT. We exhibit an intimate relationship between “reciprocity sheaves” from [7] and “modulus sheaves with transfers” from [4, 5].

CONTENTS

Introduction	1
1. Review of basic definitions and results	4
2. $\overline{\square}$ -invariance and SC-reciprocity	6
3. Relation with [7]	16
References	24

INTRODUCTION

This paper is a synthesis of [7] and [4, 5]; part of it uses results of [14] and [16].

In [7], we introduced reciprocity (pre)sheaves as a generalization of Voevodsky’s homotopy invariant (pre)sheaves with transfers, which are the main building block for constructing his triangulated categories of motives in [18]. (From now on, we shall replace *homotopy invariant* by \mathbf{A}^1 -invariant for clarity.) Let \mathbf{Sm} be the category of separated smooth schemes of finite type over k . There is an additive category \mathbf{Cor} which has the same objects as \mathbf{Sm} and whose morphisms are finite correspondences; the category \mathbf{PST} of *presheaves with transfers* is defined as the additive dual of \mathbf{Cor} [13, Lect. 1 and 2]. A presheaf with transfers F is \mathbf{A}^1 -invariant if the projection $X \times \mathbf{A}^1 \rightarrow X$ induces an isomorphism $F(X) \xrightarrow{\sim} F(X \times \mathbf{A}^1)$ for all $X \in \mathbf{Sm}$. Let $\mathbf{HI} \subset \mathbf{PST}$ be the full subcategory of \mathbf{A}^1 -invariant presheaves with transfers. The

Date: February 18, 2021.

2010 *Mathematics Subject Classification.* 19E15 (14F42, 19F15).

The first author acknowledges the support of Agence Nationale de la Recherche (ANR) under reference ANR-12-BL01-0005. The second author is supported by JSPS KAKENHI Grant (15H03606). The third author is supported by JSPS KAKENHI Grant (15K04773).

reciprocity presheaves defined in [7] form a full subcategory $\mathbf{Rec} \subset \mathbf{PST}$, which contains \mathbf{HI} .

In this paper, we introduce a new full subcategory $\mathbf{RSC} \subset \mathbf{PST}$ which is fairly close to \mathbf{Rec} and fits better with the new framework of *modulus presheaves of transfers*. The latter were introduced in [8] to construct a new triangulated category $\mathbf{MDM}^{\text{eff}}$ of motivic nature which enlarges Voevodsky's triangulated category of motives \mathbf{DM}^{eff} [13, Lect. 14]. Due to problems encountered in [8], this theory was refounded in [4, 5] and [6]. In this paper, we only use results from [4] and [5], except for the tensor structure on \mathbf{MCor} .

To give an idea of how one defines \mathbf{Rec} and \mathbf{RSC} , we need to reformulate the definition of \mathbf{A}^1 -invariance. Recall [13, Lem. 2.16] that the inclusion $\mathbf{HI} \rightarrow \mathbf{PST}$ has a left adjoint

$$(0.1) \quad h_0^{\mathbf{A}^1} : \mathbf{PST} \rightarrow \mathbf{HI}.$$

Thus $F \in \mathbf{PST}$ is in \mathbf{HI} if and only if for any $X \in \mathbf{Sm}$ and $a \in F(X)$, the map $\mathbb{Z}_{\text{tr}}(X) \rightarrow F$ in \mathbf{PST} associated to a by Yoneda's lemma factors through $h_0^{\mathbf{A}^1}(X) := h_0^{\mathbf{A}^1}(\mathbb{Z}_{\text{tr}}(X))$, where $\mathbb{Z}_{\text{tr}}(X)$ is the presheaf with transfers represented by X . To define reciprocity presheaves, we introduced in [7] bigger quotients $h(M)$ of $\mathbb{Z}_{\text{tr}}(X)$ associated to a *modulus pair* $M = (\overline{X}, X^\infty)$, consisting of a proper scheme \overline{X} over k and an effective Cartier divisor X^∞ on it, such that $X = \overline{X} \setminus |X^\infty|$. Then a presheaf with transfers $F \in \mathbf{PST}$ belongs to \mathbf{Rec} [7, Definition 2.1.3] if

(*) For any quasi-affine $X \in \mathbf{Sm}$ and any $a \in F(X)$, the associated map $a : \mathbb{Z}_{\text{tr}}(X) \rightarrow F$ factors through $h(M)$ for some M as above.

The definition of the quotients $h(M)$ is very technical; it is inspired by the theorem of Rosenlicht-Serre on reciprocity for morphisms from curves to commutative algebraic groups [17, Ch. III].

Let us now recall the story of [4, 5]. We define a category \mathbf{MCor} : its objects are modulus pairs $M = (\overline{X}, X^\infty)$ as above such that $M^\circ = \overline{X} - |X^\infty| \in \mathbf{Sm}$: this is called the *interior* of M . Morphisms of \mathbf{MCor} are finite correspondences between interiors satisfying an admissibility condition with respect to X^∞ (see Definition 1.1.1). Let \mathbf{MPST} be the additive dual of \mathbf{MCor} . There is a pair of adjoint functors

$$\mathbf{MPST} \begin{array}{c} \xrightarrow{\omega!} \\ \xleftarrow{\omega^*} \end{array} \mathbf{PST}.$$

Here ω^* is induced by the ‘‘interior’’ functor

$$\omega : \mathbf{MCor} \rightarrow \mathbf{Cor} : (\overline{X}, X^\infty) \mapsto \overline{X} \setminus |X^\infty|,$$

and $\omega_!$ is the left Kan extension of ω .

Let $\overline{\square} = (\mathbf{P}^1, \infty) \in \mathbf{MCor}$: we say that $F \in \mathbf{MPST}$ is $\overline{\square}$ -invariant if the ‘‘projection’’ $M \otimes \overline{\square} \rightarrow M$ induces an isomorphism $F(M) \xrightarrow{\sim} F(M \otimes \overline{\square})$ for all $M \in \mathbf{MCor}$ (see §1.2 for the monoidal structure \otimes on \mathbf{MCor}). We let $\mathbf{CI} \subset \mathbf{MPST}$ denote the full subcategory of $\overline{\square}$ -invariant objects.

We show in Theorem 2.1.8 that the inclusion $\mathbf{CI} \rightarrow \mathbf{MPST}$ has a left adjoint $h_0^{\overline{\square}} : \mathbf{MPST} \rightarrow \mathbf{CI}$. Define $h_0(M) \in \mathbf{PST}$ to be $\omega_! h_0^{\overline{\square}} \mathbb{Z}_{\text{tr}}(M)$, where $\mathbb{Z}_{\text{tr}}(M) \in \mathbf{MPST}$ is the presheaf represented by M and $\omega_!$ is as before. Then \mathbf{RSC} is the full subcategory of \mathbf{PST} consisting of those presheaves verifying Condition (*) above, modified by dropping the quasi-affine condition on X and replacing $h(M)$ by $h_0(M)$.

Our main results are the following.

Theorem 1.

- (1) (Corollary 2.3.4). We have $\mathbf{HI} \subset \mathbf{RSC}$.
- (2) (Th. 2.3.3 and Prop. 2.3.7). We have $\omega_!(\mathbf{CI}) = \mathbf{RSC}$. The induced functor $\omega_{\mathbf{CI}} : \mathbf{CI} \rightarrow \mathbf{RSC}$ has a fully faithful right adjoint $\omega^{\mathbf{CI}} : \mathbf{RSC} \rightarrow \mathbf{CI}$.

Theorem 2.

- (1) (Th. 3.2.1). Let $M = (\overline{X}, Y) \in \mathbf{MCor}$ be such that $X := \overline{X} \setminus |Y|$ is quasi-affine. Then $h_0(M) = h(M)$. Consequently, we have $\mathbf{RSC} \subset \mathbf{Rec}$.
- (2) (Cor. 3.2.3). We have

$$\mathbf{RSC} \cap \mathbf{NST} = \mathbf{Rec} \cap \mathbf{NST}.$$

Here, $\mathbf{NST} \subset \mathbf{PST}$ is the full subcategory of Nisnevich sheaves with transfers [18].

Voevodsky’s theory of homotopy invariant presheaves with transfers relies on an algebro-geometric version of classical homotopy theory, where the rôle of the interval is played by the affine line \mathbf{A}^1 . Reciprocity presheaves with transfers were introduced in [7] to generalize the former, based on the completely different idea of reciprocity à la Rosenlicht-Serre. Conversely, the above theorems say that one may largely understand them in terms of a more sophisticated homotopy theory, based on $\overline{\square}$ rather than \mathbf{A}^1 . This is a remarkable fact.

Remark 3. In [7, Conjecture 1 (1)], it is conjectured that the Cousin complex attached to $F \in \mathbf{Rec} \cap \mathbf{NST}$ is exact. This is proved for $F \in \mathbf{RSC} \cap \mathbf{NST}$ in [16, Cor. 3]. Thus, Theorem 2 (2) permits us to deduce the full statement of the original conjecture.

Corrections. In the first version of this paper [9], we made the following two claims about the functor $\omega_{\mathbf{CI}}$ from Theorem 1 (2): it induces 1) a monoidal structure on \mathbf{RSC} from the one on \mathbf{CI} , and 2) an equivalence of categories

$$\mathbf{CI} \cap \mathbf{MNST} \xrightarrow{\sim} \mathbf{RSC} \cap \mathbf{NST},$$

where $\mathbf{MNST} \subset \mathbf{MPST}$ is the full subcategory of ‘*Modulus Nisnevich sheaves with transfers*’ (see §1.4). Both proofs have turned out to be incorrect. The mistake in 2) originates in a false statement in the initial version of [16], which has been removed from its published version. See Remark 2.4.4 for a counterexample in characteristic zero.

Acknowledgments. Part of this work was done while the authors stayed at the university of Regensburg supported by the SFB grant “Higher Invariants”. Another part was done in a Research in trio in CIRM, Luminy. Yet another part was done while the third author was visiting IMJ-PRG supported by the Foundation Sciences Mathématiques de Paris. Yet another part was done while the first author was visiting Tohoku University and the Tokyo Institute of Technology. We are grateful to the support and hospitality received in all places.

The authors express their deep gratitude to Kay Rülling for pointing out the errors mentioned in the previous paragraph. They are also grateful to the referee for careful reading and useful comments.

Notation and conventions. Throughout this paper we work over a base field k . Denote by \mathbf{Sch} the category of separated schemes of finite type over k , and by \mathbf{Sm} the full subcategory of \mathbf{Sch} consisting of all smooth k -schemes.

1. REVIEW OF BASIC DEFINITIONS AND RESULTS

1.1. **Modulus pairs.** The following definitions (1) and (2) are taken from [4, Definitions 1.1.1, 1.3.1].

Definition 1.1.1.

- (1) A pair $M = (\overline{X}, X_\infty)$ of $\overline{X} \in \mathbf{Sch}$ and an effective Cartier divisor X_∞ on \overline{X} is called a *modulus pair* if $\overline{X} \setminus |X_\infty| \in \mathbf{Sm}$. It is called *proper* if \overline{X} is proper over k .
- (2) Let $M = (\overline{X}, X_\infty)$, $N = (\overline{Y}, Y_\infty)$ be two proper modulus pairs and put $X = \overline{X} \setminus |X_\infty|$, $Y = \overline{Y} \setminus |Y_\infty|$. We define $\mathbf{MCor}(M, N)$ to be the subgroup of $\mathbf{Cor}(X, Y)$ generated by all elementary correspondences $V \in \mathbf{Cor}(X, Y)$ such that the closure \overline{V} of V in $\overline{X} \times \overline{Y}$ satisfies $\nu^*(\overline{X} \times Y_\infty) \leq \nu^*(X_\infty \times \overline{Y})$, where $\nu : \overline{V}^N \rightarrow \overline{X} \times \overline{Y}$ is the composition of the normalization $\overline{V}^N \rightarrow \overline{V}$

and the inclusion $\overline{V} \hookrightarrow \overline{X} \times \overline{Y}$. We call these correspondences *admissible* (with respect to (M, N)). This defines a category \mathbf{MCor} of proper modulus pairs.

There is a functor

$$(1.1) \quad \omega : \mathbf{MCor} \rightarrow \mathbf{Cor}$$

defined by $\omega(\overline{X}, X_\infty) = \overline{X} \setminus |X_\infty|$.

The following is basic to his paper.

Lemma 1.1.2 ([6, §2.1]). *The assignment*

$$(\overline{X}, X_\infty) \otimes (\overline{Y}, Y_\infty) = (\overline{X} \times \overline{Y}, X_\infty \times Y_\infty + \overline{X} \times Y_\infty).$$

defines a symmetric monoidal structure on \mathbf{MCor} , with unit object $\mathbf{1} = (\text{Spec } k, \emptyset)$. The functor ω of (1.1) is symmetric monoidal.

1.2. Modulus presheaves with transfers. Here is the definition of our main object of study (see [4, Definition 2.1.1, Notation 2.1.2]).

Definition 1.2.1.

- (1) We denote by \mathbf{MPST} the abelian category of all additive functors $\mathbf{MCor}^{\text{op}} \rightarrow \mathbf{Ab}$.
- (2) For $M \in \mathbf{MCor}$, we denote by $\mathbb{Z}_{\text{tr}}(M) \in \mathbf{MPST}$ the object represented by M .

By [13, Def. 8.2] and [11, Appendix] we have the following.

Proposition 1.2.2. *The category \mathbf{MPST} has a symmetric monoidal structure that extends the tensor structure of Lemma 1.1.2 via the additive Yoneda functor. It admits an internal Hom such that*

$$(1.2) \quad \underline{\text{Hom}}_{\mathbf{MPST}}(\mathbb{Z}_{\text{tr}}(M), F)(N) = F(M \otimes N)$$

for $M, N \in \mathbf{MCor}$ and $F \in \mathbf{MPST}$.

1.3. Relation with PST. The functor ω of (1.1) induces a functor $\omega^* : \mathbf{PST} \rightarrow \mathbf{MPST}$, $\omega^*(F) = F \circ \omega$.

Proposition 1.3.1.

- (1) *The functor ω^* is fully faithful and exact.*
- (2) *There is a left adjoint $\omega_! : \mathbf{MPST} \rightarrow \mathbf{PST}$ of ω^* , which is monoidal and exact. We have*

$$\omega_! F(X) \simeq \varinjlim_{M \in \mathbf{MSm}(X)} F(M) \quad (F \in \mathbf{MPST}, X \in \mathbf{Sm}).$$

In (2), $\mathbf{MSm}(X)$ is the inverse system $\{M = (\overline{X}, X_\infty) \in \mathbf{MCor} \mid X = \overline{X} \setminus |X_\infty|\}$, where transition maps are given by the diagonal $X \subset X \times X$ whenever it defines a morphism in \mathbf{MCor} .

Proof. See [4, Prop. 2.2.1 and (2.2.1)]. The monoidality of $\omega_!$ follows that of (1.1). \square

1.4. Modulus sheaves with transfers. In [5, Lemma-Definition 4.2.1], we define a full subcategory $\mathbf{MNST} \subset \mathbf{MPST}$ of “modulus Nisnevich sheaves with transfers”. In this paper we need the following:

Proposition 1.4.1.

- (1) *The category \mathbf{MNST} is abelian; the full embedding $i_{\mathbf{Nis}} : \mathbf{MNST} \hookrightarrow \mathbf{MPST}$ has an exact left adjoint $a_{\mathbf{Nis}}$ (“sheafification”).*
- (2) *The functors $\omega_!$ and ω^* of Proposition 1.3.1 preserve \mathbf{MNST} and \mathbf{NST} ; they induce an adjunction $(\omega_{\mathbf{Nis}}, \omega^{\mathbf{Nis}})$ between these two categories, and $\omega_{\mathbf{Nis}}, \omega^{\mathbf{Nis}}$ are both exact. Moreover, the pair $(\omega_!, \omega_{\mathbf{Nis}})$ commutes with the sheafification functors $a_{\mathbf{Nis}}$ and $a_{\mathbf{Nis}}^V : \mathbf{PST} \rightarrow \mathbf{NST}$ [18, Th. 3.1.4].*

Proof. See [5, Theorem 4.2.4] for (1) and [5, Prop. 6.2.1] for (2). \square

2. $\overline{\square}$ -INVARIANCE AND SC-RECIPROCITY

2.1. $\overline{\square}$ -INVARIANCE.

Definition 2.1.1. Let $\overline{\square} = (\mathbf{P}^1, \infty)$, and write $p : \overline{\square} \rightarrow \mathbf{1}$ for the canonical morphism. We say $F \in \mathbf{MPST}$ is $\overline{\square}$ -invariant if the projection map $1_M \otimes p : M \otimes \overline{\square} \rightarrow M$ induces an isomorphism $p^* : F(M) \xrightarrow{\sim} F(M \otimes \overline{\square})$ for any $M \in \mathbf{MCor}$. We define \mathbf{CI} to be the full subcategory of \mathbf{MPST} consisting of all objects having $\overline{\square}$ -invariance.

Lemma 2.1.2. *The category \mathbf{CI} is closed under taking subobjects, quotients and extensions in \mathbf{MPST} .*

Proof. Since the zero section $i_0 : \mathbf{1} \rightarrow \overline{\square}$ is right inverse to p , $p^* : F(M) \rightarrow F(M \otimes \overline{\square})$ is an isomorphism if and only if $i_0^* : F(M \otimes \overline{\square}) \rightarrow F(M)$ is injective. This implies that $\overline{\square}$ -invariance is preserved under taking subobjects. The remaining assertions then follow by the five lemma. \square

Consider the multiplication map

$$\mu : \mathbf{A}^1 \times \mathbf{A}^1 \rightarrow \mathbf{A}^1; \quad (x, y) \mapsto (xy),$$

Let $\Gamma \subset \mathbf{A}^1 \times \mathbf{A}^1 \times \mathbf{A}^1$ be the graph of μ .

Lemma 2.1.3 ([6, Lem. 5.1.1]). *We have $\Gamma \in \mathbf{MCor}(\overline{\square} \otimes \overline{\square}, \overline{\square})$. In other words, the finite correspondence μ is admissible.*

Definition 2.1.4. For $F \in \mathbf{MPST}$, define $h_0^\square(F) \in \mathbf{MPST}$ by:

$$(2.1) \quad h_0^\square(F)(M) = \text{Coker} \left(F(M \otimes \square) \xrightarrow{i_0^* - i_1^*} F(M) \right) \quad (M \in \mathbf{MCor})$$

where i_ε^* for $\varepsilon = 0, 1$ is the pullback by the section $i_\varepsilon : \mathbf{1} \rightarrow \square$ sending $\text{Spec } k$ to $\varepsilon \in \mathbf{A}^1(k)$. For $M \in \mathbf{MCor}$, we write $h_0^\square(M) = h_0^\square(\mathbb{Z}_{\text{tr}}(M))$.

Proposition 2.1.5. *Let $F \in \mathbf{MPST}$.*

- (1) *The following conditions are equivalent.*
 - (i) $F \in \mathbf{CI}$.
 - (ii) *The natural map $F \rightarrow h_0^\square(F)$ is an isomorphism.*
 - (iii) *For any $M \in \mathbf{MCor}$ and $a \in F(M)$, the Yoneda map $\tilde{a} : \mathbb{Z}_{\text{tr}}(M) \rightarrow F$ factors through $h_0^\square(M)$.*
- (2) *We have $h_0^\square(F) \in \mathbf{CI}$ and the induced functor*

$$h_0^\square : \mathbf{MPST} \rightarrow \mathbf{CI}; \quad F \mapsto h_0^\square(F)$$

gives a left adjoint of the inclusion $i^\square : \mathbf{CI} \hookrightarrow \mathbf{MPST}$.

- (3) *For any $M \in \mathbf{MCor}$, the morphism $h_0^\square(1_M \otimes p) : h_0^\square(M \otimes \square) \rightarrow h_0^\square(M)$ is an isomorphism.*

Proof. It essentially reproduces the proof of the same facts for \mathbf{A}^1 -invariant presheaves, by adding modulus. The main point is Lemma 2.1.3.

Assume (i) and take $M \in \mathbf{MCor}$. The assumption implies that $i_\varepsilon^* : F(M \otimes \square) \rightarrow F(M)$ for $\varepsilon = 0, 1$ are both inverse to $p^* : F(M) \xrightarrow{\sim} F(M \otimes \square)$ so that $i_0^* - i_1^* = 0$, which implies (ii).

Assume (ii). By (2.1) this implies that for any $M \in \mathbf{MCor}$ we have

$$(2.2) \quad i_0^* = i_1^* : F(M \otimes \square) \rightarrow F(M).$$

By Lemma 2.1.3, we have a commutative diagram

$$\begin{array}{ccc} F(M \otimes \square) & \xrightarrow{(1_M \otimes i_0)^*} & F(M) \\ (1_M \otimes \mu)^* \downarrow & & \downarrow p^* \\ F(M \otimes \square \otimes \square) & \xrightarrow{(1_{M \otimes \square} \otimes i_0)^*} & F(M \otimes \square). \end{array}$$

By this diagram and (2.2), we get

$$\begin{aligned} p^*(1_M \otimes i_0)^* &= (1_{M \otimes \square} \otimes i_0)^* \circ (1_M \otimes \mu)^* \\ &= (1_{M \otimes \square} \otimes i_1)^* \circ (1_M \otimes \mu)^* = 1_{M \otimes \square}^*. \end{aligned}$$

This proves the surjectivity of p^* , hence (i) holds. Thus (i) \iff (ii).

By the definition of $h_0^\square(F)$, for any $M \in \mathbf{MCor}$, the map

$$h_0^\square(F)(M \otimes \square) \xrightarrow{i_0^* - i_1^*} h_0^\square(F)(M)$$

is the zero map so that $h_0^\square(F)(M) \simeq h_0^\square(h_0^\square(F))(M)$. Hence the first assertion of (2) follows from the implication (ii) \Rightarrow (i).

Any $\tilde{a} : \mathbb{Z}_{\text{tr}}(M) \rightarrow F$ induces a morphism $h_0^\square(M) \rightarrow h_0^\square(F)$ which commutes with the natural transformation of (ii). Hence (iii) follows from (ii).

If (iii) holds, F is a quotient of a direct sum of $h_0^\square(M)$'s for $M \in \mathbf{MCor}$. Hence (i) holds by the first assertion of (2) and Lemma 2.1.2. This completes the proof of (1).

To show the second assertion of (2), note that (1) implies that for $F \in \mathbf{CI}$ and $M \in \mathbf{MCor}$, the natural map $\mathbb{Z}_{\text{tr}}(M) \rightarrow h_0^\square(M)$ induces an isomorphism

$$(2.3) \quad \text{Hom}_{\mathbf{MPST}}(h_0^\square(M), F) \simeq \text{Hom}_{\mathbf{MPST}}(\mathbb{Z}_{\text{tr}}(M), F).$$

For $G \in \mathbf{MPST}$, take a resolution of G of the form

$$P_1 \rightarrow P_0 \rightarrow G \rightarrow 0 \quad \text{in } \mathbf{MPST},$$

where P_1, P_0 are direct sums of representable objects. By its definition (2.1), the endofunctor h_0^\square of \mathbf{MPST} is right exact so that the above sequence induce an exact sequence

$$h_0^\square(P_1) \rightarrow h_0^\square(P_0) \rightarrow h_0^\square(G) \rightarrow 0.$$

Moreover h_0^\square commutes with direct sums again by (2.1). In view of (2.3), we conclude that the natural map $G \rightarrow h_0^\square(G)$ induces an isomorphism

$$\text{Hom}_{\mathbf{MPST}}(h_0^\square(G), F) \simeq \text{Hom}_{\mathbf{MPST}}(G, F).$$

which implies the desired claim.

It remains to prove (3). Since $h_0^\square(1_M \otimes i_0)$ is a right inverse of $h_0^\square(1_M \otimes p)$, it suffices to show that it is also a left inverse. Let $N \in \mathbf{MCor}$. For $\varphi \in \mathbf{MCor}(N, M \otimes \square)$, define

$$\tilde{\varphi} = (1_M \otimes \mu) \circ (\varphi \otimes 1_\square) \in \mathbf{MCor}(N \otimes \square, M \otimes \square).$$

Using the identities $\mu \circ (1_\square \otimes i_0) = i_0 p$ and $\mu \circ (1_\square \otimes i_1) = 1_\square$, we get

$$(i_0^* - i_1^*)\tilde{\varphi} := \tilde{\varphi} \circ (1_N \otimes i_0) - \tilde{\varphi} \circ (1_N \otimes i_1) = (1_M \otimes i_0 p) \circ \varphi - \varphi$$

which shows that $\tilde{\varphi} = h_0^\square(1_M \otimes i_0)h_0^\square(1_M \otimes p)\tilde{\varphi}$, where $\tilde{\varphi}$ is the image of φ in $h_0^\square(M \otimes \square)(N)$. \square

Definition 2.1.6. For $F \in \mathbf{MPST}$, define $h_{\square}^0(F) \in \mathbf{MPST}$ by

$$(2.4) \quad h_{\square}^0(F)(M) = \mathrm{Hom}_{\mathbf{MPST}}(h_0^{\square}(M), F) \quad (M \in \mathbf{MCor}).$$

Lemma 2.1.7. For $F \in \mathbf{MPST}$, $h_{\square}^0(F)$ is the maximal \square -invariant subobject of F . The induced functor

$$h_{\square}^0 : \mathbf{MPST} \rightarrow \mathbf{CI}; F \mapsto h_{\square}^0(F)$$

gives a right adjoint of the inclusion $i^{\square} : \mathbf{CI} \hookrightarrow \mathbf{MPST}$.

Proof. The fact that $h_{\square}^0(F)$ is a subobject of F follows from (2.4) and the fact that $h_0^{\square}(M)$ is a quotient of $\mathbb{Z}_{\mathrm{tr}}(M)$. The fact that $h_{\square}^0(F) \in \mathbf{CI}$ follows from Proposition 2.1.5 (3). Now let $G \subset F$ be a subobject which is in \mathbf{CI} . For $a \in F(M)$ with $M \in \mathbf{MCor}$, let $\tilde{a} : \mathbb{Z}_{\mathrm{tr}}(M) \rightarrow F$ be the corresponding map in \mathbf{MPST} . If $a \in G(M)$, \tilde{a} factors through G and hence factors through $h_0^{\square}(M)$ by Proposition 2.1.5 (1). Hence $a \in h_{\square}^0(F)(M)$ by (2.4). This proves $G \subset h_{\square}^0(F)$, which completes the proof of the first assertion. The second assertion follows easily from the first and Lemma 2.1.2. \square

Theorem 2.1.8. The category \mathbf{CI} is a Serre subcategory of \mathbf{MPST} , and is Grothendieck. The inclusion $i^{\square} : \mathbf{CI} \hookrightarrow \mathbf{MPST}$ has

- (i) a left adjoint given by $F \mapsto h_0^{\square}(F)$;
- (ii) a right adjoint given by $F \mapsto h_{\square}^0(F)$, where

$$h_{\square}^0 F(M) = \mathrm{Hom}(h_0^{\square}(M), F) \quad (M \in \mathbf{MCor}).$$

The unit (resp. counit) morphism $F \rightarrow h_0^{\square}(F)$ (resp. $h_{\square}^0(F) \rightarrow F$) is epi (resp. mono).

Proof. Everything follows from what we have proven so far, except for the Grothendieckness of \mathbf{CI} , that we want to deduce from that of \mathbf{MPST} . We cannot quite apply [4, Th. A.10.1 d)], because h_0^{\square} is not exact: it provides generators and infinite direct sums, but not their exactness. But the latter holds because i^{\square} reflects infinite direct sums (if (F_{α}) is a family of objects in \mathbf{CI} , their direct sum in \mathbf{MPST} belongs to \mathbf{CI}). \square

Proposition 2.1.9.

- (1) One has $\underline{\mathrm{Hom}}_{\mathbf{MPST}}(G, H) \in \mathbf{CI}$ for any $G \in \mathbf{MPST}$ and any $H \in \mathbf{CI}$.
- (2) Via h_0^{\square} , the symmetric monoidal structure on \mathbf{MPST} from Proposition 1.2.2 induces a symmetric monoidal structure on \mathbf{CI} by the formula

$$F \otimes_{\mathbf{CI}} G = h_0^{\square}(F \otimes_{\mathbf{MPST}} G) \quad (F, G \in \mathbf{CI}).$$

This tensor product commutes with all representable colimits.

Proof. (1) follows from (1.2) and the isomorphisms (easily checked by means of Yoneda's lemma):

$$\begin{aligned} \underline{\mathrm{Hom}}_{\mathrm{MPST}}(\overline{\square}, \underline{\mathrm{Hom}}_{\mathrm{MPST}}(G, H)) &\simeq \underline{\mathrm{Hom}}_{\mathrm{MPST}}(\overline{\square} \otimes_{\mathrm{MPST}} G, H) \\ &\simeq \underline{\mathrm{Hom}}_{\mathrm{MPST}}(G, \underline{\mathrm{Hom}}_{\mathrm{MPST}}(\overline{\square}, H)) \xleftarrow{\sim} \underline{\mathrm{Hom}}_{\mathrm{MPST}}(G, H) \end{aligned}$$

where the last isomorphism, induced by p , follows from the assumption $H \in \mathbf{CI}$.

(2) Since $h_0^{\overline{\square}}$ is a localisation by the full faithfulness of $i^{\overline{\square}}$, we have to show the following.

Claim 2.1.10. If $f \in \mathrm{Hom}_{\mathrm{MPST}}(G_1, G_2)$ is such that $h_0^{\overline{\square}}(f)$ is an isomorphism, then $g = h_0^{\overline{\square}}(f \otimes_{\mathrm{MPST}} 1_{G'})$ is an isomorphism for any $G' \in \mathrm{MPST}$.

By (co)Yoneda, it suffices to show that

$$g^* : \mathrm{Hom}_{\mathrm{MPST}}(h_0^{\overline{\square}}(G_2 \otimes_{\mathrm{MPST}} G'), H) \rightarrow \mathrm{Hom}_{\mathrm{MPST}}(h_0^{\overline{\square}}(G_1 \otimes_{\mathrm{MPST}} G'), H)$$

is an isomorphism for any $H \in \mathbf{CI}$. By adjunction, it suffices to show that

$$f^* : \mathrm{Hom}_{\mathrm{MPST}}(G_2, \underline{\mathrm{Hom}}_{\mathrm{MPST}}(G', H)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{MPST}}(G_1, \underline{\mathrm{Hom}}_{\mathrm{MPST}}(G', H))$$

which follows from (1).

Finally, the commutation of $\otimes_{\mathbf{CI}}$ with colimits follows from that of \otimes_{MPST} and $h_0^{\overline{\square}}$. \square

2.2. SC-reciprocity.

Definition 2.2.1. For $F \in \mathrm{MPST}$, we define

$$h_0(F) := \omega_! h_0^{\overline{\square}}(F) \in \mathbf{PST}.$$

For $M \in \mathbf{MCor}$, we put $h_0(M) := h_0(\mathbb{Z}_{\mathrm{tr}}(M))$.

Lemma 2.2.2. Let $M = (\overline{M}, M_\infty) \in \mathbf{MCor}$ with $M^\circ = M - |M_\infty|$. For $S \in \mathbf{Sm}$, we have

$$h_0(M)(S) = \mathrm{Coker}(i_0^* - i_1^* : \underline{\mathbf{MCor}}(\overline{\square} \otimes S, M) \rightarrow \mathbf{Cor}(S, M^\circ)),$$

where $\underline{\mathbf{MCor}}(\overline{\square} \otimes S, M)$ is the subgroup of $\mathbf{Cor}(\mathbf{A}^1 \times S, M^\circ)$ generated by all elementary correspondences Z such that

$$\varphi_Z^*(\mathbf{P}^1 \times S \times M_\infty) \leq \varphi_Z^*(\infty \times S \times \overline{M}),$$

where $\varphi_Z : \overline{Z}^{\mathbf{N}} \rightarrow \overline{Z} \hookrightarrow \mathbf{P}^1 \times S \times \overline{M}$ denotes the normalization of the closure \overline{Z} of Z in $\mathbf{P}^1 \times S \times \overline{M}$.

Proof. This follows from Definition 2.1.4 and Proposition 1.3.1 (2). \square

Remark 2.2.3. Lemma 2.2.2 implies isomorphisms

$$h_0(M)(\mathrm{Spec} k) \simeq H_0^S(\overline{M}, M_\infty) \simeq \mathrm{CH}_0(M),$$

where $H_n^S(\overline{M}, M_\infty)$ for $n \in \mathbb{Z}$ is the *Suslin homology* considered in [15, Definition 3.1] and $\mathrm{CH}_0(M) = \mathrm{CH}_0(\overline{M}|M_\infty)$ is the Chow group with modulus considered in [12].

Definition 2.2.4.

- (1) Let $F \in \mathbf{PST}$, $X \in \mathbf{Sm}$ and $a \in F(X) = \mathrm{Hom}_{\mathbf{PST}}(\mathbb{Z}_{\mathrm{tr}}(X), F)$. We say $M = (\overline{X}, X_\infty) \in \mathbf{MCor}$ is an *SC-modulus* for a if $X = \overline{X} \setminus |X_\infty|$ and $a : \mathbb{Z}_{\mathrm{tr}}(X) \rightarrow F$ factors through $\mathbb{Z}_{\mathrm{tr}}(X) \rightarrow h_0(M)$. (SC stands for ‘‘Suslin complex’’.)
- (2) We say $F \in \mathbf{PST}$ has *SC-reciprocity* if, for any $X \in \mathbf{Sm}$, any $a \in F(X)$ has an SC-modulus $M \in \mathbf{MSm}(X)$.
- (3) We define **RSC** to be the full subcategory of **PST** consisting of all objects having SC-reciprocity.

Remark 2.2.5. The category **RSC** is closed under subobjects and quotient objects in **PST**. This is obvious from the definition. In particular, **RSC** is abelian and the inclusion functor $i^\natural : \mathbf{RSC} \rightarrow \mathbf{PST}$ is exact.

Recall that $i^{\square} : \mathbf{CI} \hookrightarrow \mathbf{MPST}$ denotes the inclusion.

Proposition 2.2.6. *The functor $\rho := \omega_! i^{\square} h_{\square}^0 \omega^*$ sends **PST** into **RSC**, and is right adjoint to the inclusion $i^\natural : \mathbf{RSC} \hookrightarrow \mathbf{PST}$.*

Proof. For $F \in \mathbf{PST}$ and $X \in \mathbf{Sm}$, we have by successive adjunctions and by Proposition 1.3.1 (2):

$$\begin{aligned} (2.5) \quad \rho F(X) &= \varinjlim_{M \in \mathbf{MSm}(X)} i^{\square} h_{\square}^0 \omega^* F(M) \\ &= \varinjlim_{M \in \mathbf{MSm}(X)} \mathbf{MPST}(h_0^{\square}(M), \omega^* F) = \varinjlim_{M \in \mathbf{MSm}(X)} \mathbf{PST}(h_0(M), F) \end{aligned}$$

which realises ρF as the largest subobject of F which is in **RSC**. \square

Corollary 2.2.7. *Let $F \in \mathbf{PST}$. The counit map*

$$(2.6) \quad i^\natural \rho F \rightarrow F$$

of the adjunction in Proposition 2.2.6 agrees with the counit map of the adjunction $(\omega_! i^{\square}, h_{\square}^0 \omega^)$. Moreover, $F \in \mathbf{RSC}$ if and only if (2.6) is an isomorphism.*

Proof. The first statement simply restates the computation in (2.5). The second one follows from Proposition 2.2.6 and the definition of SC-reciprocity. \square

Question 2.2.8. Is **RSC** closed under extensions in **PST**?

This question appears to be very difficult (compare [7, Question 1]). We can only offer a trivial reduction:

Proposition 2.2.9. *For $F \in \mathbf{PST}$, the following are equivalent:*

(i) *There exist $G, H \in \mathbf{RSC}$ and an exact sequence in \mathbf{PST}*

$$(2.7) \quad 0 \rightarrow G \rightarrow F \rightarrow H \rightarrow 0.$$

(ii) *The cokernel of (2.6) is in \mathbf{RSC} .*

Proof. (ii) \Rightarrow (i) is obvious. Conversely, assume (i). Applying the left exact functor ρ , we get an exact sequence in \mathbf{RSC} (defining C)

$$(2.8) \quad 0 \rightarrow G \rightarrow \rho F \rightarrow H \rightarrow C \rightarrow 0.$$

The counit map (2.6) sends (2.8) to (2.7). A diagram chase then gives us the exact sequence in \mathbf{PST}

$$0 \rightarrow \rho F \rightarrow F \rightarrow C \rightarrow 0$$

which concludes the proof. \square

2.3. Relations between \mathbf{CI} , \mathbf{HI} and \mathbf{RSC} . Recall that $\mathbf{HI} \subset \mathbf{PST}$ is the full subcategory of \mathbf{A}^1 -invariant presheaves with transfers.

Lemma 2.3.1. *For $H \in \mathbf{PST}$, $H \in \mathbf{HI}$ if and only if $\omega^*H \in \mathbf{CI}$.*

Proof. This follows from the fact that for $M = (\overline{M}, M_\infty) \in \mathbf{MCor}$ with $M^\circ = \overline{M} - M_\infty$, we have

$$\omega^*F(M) = F(M^\circ) \quad \text{and} \quad \omega^*F(M \otimes \overline{\square}) = F(M^\circ \times \mathbf{A}^1).$$

\square

Proposition 2.3.2. *The composite $\mathbf{MPST} \xrightarrow{\omega_1} \mathbf{PST} \xrightarrow{h_0^{\mathbf{A}^1}} \mathbf{HI}$ factors through $\mathbf{MPST} \xrightarrow{h_0^{\overline{\square}}} \mathbf{CI}$, inducing a functor $\omega_h : \mathbf{CI} \rightarrow \mathbf{HI}$. This functor is right exact and monoidal for the \otimes -structures given on \mathbf{CI} by Theorem 2.1.8 (2), and analogously on \mathbf{HI} . It has an exact right adjoint ω^h , given by the restriction of ω^* to \mathbf{HI} .*

Proof. The first claim and the monoidality of ω_h follow from that of ω_1 , as $\omega_1 \mathbb{Z}_{\text{tr}}(\overline{\square}) = \mathbb{Z}_{\text{tr}}(\mathbf{A}^1)$. The existence, characterisation and exactness of ω^h follows from Lemma 2.3.1, and the right exactness of ω_h then follows. \square

Theorem 2.3.3. *If $F \in \mathbf{CI}$, then $\omega_1 F \in \mathbf{RSC}$.*

Proof. We have a commutative diagram, for any $F \in \mathbf{MPST}$:

$$(2.9) \quad \begin{array}{ccccc} \omega_! i^{\square} h_{\square}^0 F & \xrightarrow[\text{(a)}]{\omega_! i^{\square} h_{\square}^0 \eta_F} & \omega_! i^{\square} h_{\square}^0 \omega^* \omega_! F & & \\ \omega_! \varepsilon'_F \downarrow \text{(c)} & & \omega_! \varepsilon'_{\omega^* \omega_! F} \downarrow \text{(d)} & & \\ \omega_! F & \xrightarrow[\text{(b)}]{\omega_! \eta_F} & \omega_! \omega^* \omega_! F & \xrightarrow[\text{(e)}]{\varepsilon_{\omega_! F}} & \omega_! F. \end{array}$$

Here, η and ε are the unit and counit of the adjunction $(\omega_!, \omega^*)$, while ε' is the counit of the adjunction $(i^{\square}, h_{\square}^0)$. We have $(e) \circ (b) = 1_{\omega_! F}$ by the adjunction identities; since ω^* is fully faithful, (e) is an isomorphism hence so is (b). This shows that (c) factors through (a). On the other hand, ε' is mono by Theorem 2.1.8, hence so are (c) and (d) since $\omega_!$ is exact. Finally, the diagram boils down to two successive monomorphisms

$$(2.10) \quad \omega_! i^{\square} h_{\square}^0 F \hookrightarrow i^{\sharp} \rho \omega_! F \hookrightarrow \omega_! F$$

with composition $\omega_! \varepsilon'_F$. Therefore, $F \in \mathbf{CI} \Rightarrow \omega_! F \in \mathbf{RSC}$. \square

Corollary 2.3.4. *We have $\mathbf{HI} \subset \mathbf{RSC}$.*

Proof. Let $F \in \mathbf{HI}$. By Lemma 2.3.1, $\omega^* F \in \mathbf{CI}$, hence

$$F \simeq \omega_! \omega^* F \in \mathbf{RSC}$$

by Theorem 2.3.3. (See [16, Lemma 1.22] for a simpler proof.) \square

Corollary 2.3.5. *For any $F \in \mathbf{MPST}$, $h_0(F) \in \mathbf{RSC}$.*

Proof. This follows from Proposition 2.1.5 and Theorem 2.3.3. \square

Corollary 2.3.6. *The inclusion functor $i^{\sharp} : \mathbf{RSC} \hookrightarrow \mathbf{PST}$ has a pro-left adjoint ℓ .*

Proof. It suffices to show that ℓ is defined on the generators $\mathbb{Z}_{\text{tr}}(X)$. Since $h_0(M) \in \mathbf{RSC}$ for any $M \in \mathbf{MSm}(X)$ by Corollary 2.3.5, we have $\ell \mathbb{Z}_{\text{tr}}(X) = \varprojlim_{M \in \mathbf{MSm}(X)} h_0(M)$. \square

Proposition 2.3.7. *There exist unique functors $\omega_{\mathbf{CI}}$ and $\omega^{\mathbf{CI}}$ that make the two diagrams*

$$\begin{array}{ccc} \mathbf{CI} & \xrightarrow{i^{\square}} & \mathbf{MPST} \\ \omega_{\mathbf{CI}} \downarrow & & \omega_! \downarrow \\ \mathbf{RSC} & \xrightarrow{i^{\sharp}} & \mathbf{PST} \end{array} \quad \begin{array}{ccc} \mathbf{CI} & \xleftarrow{h_{\square}^0} & \mathbf{MPST} \\ \omega^{\mathbf{CI}} \uparrow & & \omega^* \uparrow \\ \mathbf{RSC} & \xrightarrow{i^{\sharp}} & \mathbf{PST} \end{array}$$

commutative, where i^\sharp is the inclusion. Moreover, $\omega^{\mathbf{CI}}$ is right adjoint to $\omega_{\mathbf{CI}}$. The counit map $\varepsilon : \omega_{\mathbf{CI}}\omega^{\mathbf{CI}} \Rightarrow \mathrm{Id}_{\mathbf{RSC}}$ is an isomorphism, $\omega_{\mathbf{CI}}$ is a localisation (in particular, is essentially surjective) and $\omega^{\mathbf{CI}}$ is fully faithful. Finally, $\omega_{\mathbf{CI}}$ is exact and $\omega^{\mathbf{CI}}$ is left exact.

Proof. The existence of $\omega_{\mathbf{CI}}$ is the contents of Theorem 2.3.3, and $\omega^{\mathbf{CI}}$ is defined by the commutativity of the diagram. For the second assertion, let $F \in \mathbf{CI}$ and $G \in \mathbf{RSC}$. Using two successive adjunctions, we compute:

$$\begin{aligned} \mathbf{CI}(F, \omega^{\mathbf{CI}}G) &= \mathbf{CI}(F, h_{\square}^0 \omega^* i^\sharp G) \simeq \mathbf{PST}(\omega_! i^{\square} F, i^\sharp G) \\ &= \mathbf{PST}(i^\sharp \omega_{\mathbf{CI}} F, i^\sharp G) \simeq \mathbf{RSC}(\omega_{\mathbf{CI}} F, G) \end{aligned}$$

where the last isomorphism uses the (tautological) full faithfulness of i^\sharp . So the adjunction $(\omega_{\mathbf{CI}}, \omega^{\mathbf{CI}})$ is obtained by ‘‘cancelling’’ i^\sharp from the adjunction $(\omega_! i^{\square}, h_{\square}^0 \omega^*)$, after applying Theorem 2.3.3. Therefore the third assertion follows from Corollary 2.2.7, and the next two are standard consequences [4, Lemma A.3.1]. The exactness of $\omega_{\mathbf{CI}}$ follows from the exactness of i^{\square} and $\omega_!$ (as well as the full faithfulness of i^\sharp), and $\omega^{\mathbf{CI}}$ is left exact as a right adjoint. \square

Corollary 2.3.8. *The category \mathbf{RSC} is Grothendieck.*

Proof. This follows from the same fact for \mathbf{CI} (Theorem 2.1.8), the adjunction $(\omega_{\mathbf{CI}}, \omega^{\mathbf{CI}})$ and [4, Th. A.10.1 d)]. \square

Proposition 2.3.9. *Let*

$$h_0^{\mathrm{rec}} : \mathbf{RSC} \rightarrow \mathbf{HI}$$

be the restriction of $h_0^{\mathbf{A}^1} : \mathbf{PST} \rightarrow \mathbf{HI}$ from (0.1). Then h_0^{rec} is a left adjoint of the inclusion $\mathbf{HI} \hookrightarrow \mathbf{RSC}$ from Corollary 2.3.4. We have a natural isomorphism $\omega_h \simeq h_0^{\mathrm{rec}} \omega_{\mathbf{CI}}$ (see Proposition 2.3.2 for ω_h).

Proof. The first claim follows immediately from the fact that $h_0^{\mathbf{A}^1}$ is a left adjoint to the inclusion $\mathbf{HI} \hookrightarrow \mathbf{PST}$. To show the second, we apply the natural isomorphism $\omega_h h_0^{\square} G \simeq h_0^{\mathbf{A}^1} \omega_! G$ from Proposition 2.3.2 to $G = i^{\square} F$ for $F \in \mathbf{CI}$ to get a natural isomorphism

$$\omega_h F \simeq \omega_h h_0^{\square} i^{\square} F \simeq h_0^{\mathbf{A}^1} \omega_! i^{\square} F \simeq h_0^{\mathbf{A}^1} i^\sharp \omega_{\mathbf{CI}} F \simeq h_0^{\mathrm{rec}} \omega_{\mathbf{CI}} F$$

as requested. \square

2.4. Sheaves in \mathbf{RSC} . Let $\mathbf{NST} \subset \mathbf{PST}$ be the full subcategory of Nisnevich sheaves with transfers [18, Th. 3.1.4]. Recall that the objects of \mathbf{NST} are those $F \in \mathbf{PST}$ whose restriction F_X to X_{Nis} is a sheaf for any $X \in \mathbf{Sm}$, where X_{Nis} denotes the small Nisnevich

site of X . By [18, Th. 3.1.4] the inclusion $i_{\text{Nis}}^V : \mathbf{NST} \rightarrow \mathbf{PST}$ has an exact left adjoint a_{Nis}^V such that for any $F \in \mathbf{PST}$ and $X \in \mathbf{Sm}$, $(a_{\text{Nis}}^V F)_X$ is the Nisnevich sheafification of F_X as a presheaf on X_{Nis} . Let $\mathbf{RSC}_{\text{Nis}} = \mathbf{RSC} \cap \mathbf{NST}$ and $\mathbf{CI}_{\text{Nis}} = \mathbf{CI} \cap \mathbf{MNST}$ (see §1.4 for \mathbf{MNST}). We admit the following theorem.

Theorem 2.4.1. *Assume k is perfect. Write*

$$\mathbf{CI}^{sp} = \{F \in \mathbf{CI} \mid \text{the unit map } F \rightarrow \omega^{\mathbf{CI}} \omega_{\mathbf{CI}} F \text{ is injective.}\}$$

- (1) [16, Th. 0.1 and 0.4] *One has $a_{\text{Nis}}^V(\mathbf{RSC}) = \mathbf{RSC}_{\text{Nis}}$ and $a_{\text{Nis}} \mathbf{CI}^{sp} \subset \mathbf{CI}_{\text{Nis}}$. (See Proposition 1.4.1 (1) for a_{Nis} .)*
- (2) [14, Cor. 4.16]. *One has $\omega^{\mathbf{CI}}(\mathbf{RSC}_{\text{Nis}}) \subset \mathbf{CI}_{\text{Nis}}$.*

Corollary 2.4.2. *The category $\mathbf{RSC}_{\text{Nis}}$ is Grothendieck.*

Proof. Since a_{Nis}^V is exact, so is its restriction to \mathbf{RSC} . The corollary now follows from Corollary 2.3.8 and (again) [4, Th. A.10.1 d)]. \square

Theorem 2.4.3. *Assume k is perfect.*

- (1) *The functor ρ of Proposition 2.2.6 sends \mathbf{NST} into $\mathbf{RSC}_{\text{Nis}}$. It yields a right adjoint ρ_{Nis} to the inclusion $i_{\text{Nis}}^{\natural} : \mathbf{RSC}_{\text{Nis}} \hookrightarrow \mathbf{NST}$.*
- (2) *The functor $\omega_{\mathbf{CI}}$ of Proposition 2.3.7 sends \mathbf{CI}_{Nis} to $\mathbf{RSC}_{\text{Nis}}$. The induced functor $\omega_{\mathbf{CI}}^{\text{Nis}} : \mathbf{CI}_{\text{Nis}} \rightarrow \mathbf{RSC}_{\text{Nis}}$ is left adjoint to the fully faithful functor $\omega_{\text{Nis}}^{\mathbf{CI}} : \mathbf{RSC}_{\text{Nis}} \rightarrow \mathbf{CI}_{\text{Nis}}$ given by Theorem 2.4.1 (2). Moreover, there is a natural isomorphism*

$$(2.11) \quad a_{\text{Nis}}^V \omega_{\mathbf{CI}} F \simeq \omega_{\mathbf{CI}}^{\text{Nis}} a_{\text{Nis}} F$$

for any $F \in \mathbf{CI}^{sp}$.

Proof. Let $F \in \mathbf{NST}$. Considering F as an object of \mathbf{PST} , we may view ρF as the largest subobject of F which belongs to \mathbf{RSC} (see Proposition 2.2.6). Applying the left exact functor a_{Nis}^V to this inclusion, we get a sequence

$$\rho F \rightarrow a_{\text{Nis}}^V \rho F \rightarrow a_{\text{Nis}}^V F = F,$$

where the second map is a monomorphism. But the middle term is in \mathbf{RSC} by Theorem 2.4.1 (1). Hence the first map must be an isomorphism, which implies the first claim of (1). The last claim now follows easily from the adjunction in Proposition 2.2.6.

The first assertion of (2) is obvious since $\omega_{\mathbf{CI}}$ preserves Nisnevich sheaves by Proposition 1.4.1. The second one then follows easily from Proposition 2.3.7. Given the natural isomorphism $\omega_{\mathbf{CI}}^{\text{Nis}} \simeq i_{\text{Nis}} \omega_{\text{Nis}}^{\mathbf{CI}}$, this implies the last assertion by taking left adjoints. \square

Remark 2.4.4. The functor $\omega_{\mathbf{CI}}^{\text{Nis}}$ is not conservative. Assume $\text{ch}(k) = 0$. Let $F \in \mathbf{CI}$ be the image of the unit map

$$h_0^{\square}(\mathbf{P}^1, 2\infty) \rightarrow \omega^{\mathbf{CI}} \omega_{\mathbf{CI}} h_0^{\square}(\mathbf{P}^1, 2\infty).$$

Then $F \in \mathbf{CI}^{sp}$, hence $a_{\text{Nis}}F \in \mathbf{CI}$ by Theorem 2.4.1 (1). We claim that the unit map $\iota : a_{\text{Nis}}F \rightarrow \omega_{\text{Nis}}^{\mathbf{CI}} \omega_{\mathbf{CI}}^{\text{Nis}} a_{\text{Nis}}F$ is not surjective. To see this, first note that by the exactness of $\omega_{\mathbf{CI}}$, we have

$$h_0(\mathbf{P}^1, 2\infty) \rightarrow \omega_{\mathbf{CI}}F \hookrightarrow \omega_{\mathbf{CI}} \omega^{\mathbf{CI}} h_0(\mathbf{P}^1, 2\infty) = h_0(\mathbf{P}^1, 2\infty),$$

and hence $\omega_{\mathbf{CI}}F \cong h_0(\mathbf{P}^1, 2\infty)$. Then by (2.11) and [15, Thm. 1.1], we have isomorphisms

$$\omega_{\mathbf{CI}}^{\text{Nis}} a_{\text{Nis}}F \simeq a_{\text{Nis}}^V \omega_{\mathbf{CI}}F \simeq \underline{\text{Pic}}(\mathbf{P}^1, 2\infty) \simeq \mathbb{Z} \oplus \mathbf{G}_a.$$

Take $(X, D) \in \mathbf{MCor}$ such that $X, D \in \mathbf{Sm}$, with X connected. Then it follows from [14, Th. 6.4] that

$$\omega_{\text{Nis}}^{\mathbf{CI}} \omega_{\mathbf{CI}}^{\text{Nis}} a_{\text{Nis}}F(X, 2mD) \simeq \mathbb{Z} \oplus H^0(X, \mathcal{O}_X((2m-1)D))$$

for any integer $m > 0$. On the other hand, one can show

$$a_{\text{Nis}}F(X, 2mD) \simeq \mathbb{Z} \oplus H^0(X, \mathcal{O}_X(mD)).$$

This implies that $G := \text{Coker}(\iota) \in \mathbf{CI}_{\text{Nis}}$ is non-zero but $\omega_{\mathbf{CI}}^{\text{Nis}}(G) = 0$.

3. RELATION WITH [7]

3.1. Review of reciprocity presheaves with transfers. In [7, Definition 2.1.3], we defined a full subcategory **Rec** of **PST**, which we now recall.

Let $(\overline{X}, Y) \in \mathbf{MCor}$ and suppose that $X = \overline{X} \setminus |Y|$ is quasi-affine. For $S \in \mathbf{Sm}$, let $\mathcal{C}_{(\overline{X}, Y)}(S)$ be the class of all finite morphisms $\varphi : \overline{C} \rightarrow \overline{X} \times S$ satisfying the following conditions:

- $\overline{C} \in \mathbf{Sch}$ is integral and normal.
- There is a generic point η of S such that $\dim \overline{C} \times_S \eta = 1$.
- The image of $\gamma_\varphi := \text{pr} \circ \varphi$ is not contained in $|Y|$, where $\text{pr} : \overline{X} \times S \rightarrow \overline{X}$ is the projection map.

For an effective Cartier divisor D on \overline{C} , we set

$$(3.1) \quad G(\overline{C}, D) := \bigcap_{x \in D} \text{Ker}(\mathcal{O}_{\overline{C}, x}^\times \rightarrow \mathcal{O}_{D, x}^\times).$$

We then define

$$\Phi(\overline{X}, Y)(S) = \bigoplus_{(\varphi : \overline{C} \rightarrow \overline{X} \times S) \in \mathcal{C}_{(\overline{X}, Y)}(S)} G(\overline{C}, \gamma_\varphi^* Y)$$

It is proved in [7, Proposition 2.2.2] that $\Phi(\overline{X}, Y)$ defines a presheaf with transfers. It is also shown there that one has $\varphi_*(\operatorname{div}_{\overline{C}}(f)) \in \mathbf{Cor}(S, X)$ for any $(\varphi : \overline{C} \rightarrow \overline{X} \times S) \in \mathcal{C}_{(\overline{X}, Y)}(S)$ and $f \in G(\overline{C}, \gamma_\varphi^* Y)$, yielding a map $\tau : \Phi(\overline{X}, Y) \rightarrow \mathbb{Z}_{\operatorname{tr}}(X)$ in \mathbf{PST} . We define

$$h(M) := \operatorname{Coker}(\tau : \Phi(\overline{X}, Y) \rightarrow \mathbb{Z}_{\operatorname{tr}}(X)) \in \mathbf{PST}.$$

Definition 3.1.1 ([7, Definition 2.1.2, Remark 2.1.6]). We say $F \in \mathbf{PST}$ has reciprocity if for any quasi-affine $X \in \mathbf{Sm}$ and $a \in F(X) = \operatorname{Hom}_{\mathbf{PST}}(\mathbb{Z}_{\operatorname{tr}}(X), F)$, there is an $M = (\overline{X}, X_\infty) \in \mathbf{MCor}$ such that $X = \overline{X} \setminus |X_\infty|$ and $a : \mathbb{Z}_{\operatorname{tr}}(X) \rightarrow F$ factors through $\mathbb{Z}_{\operatorname{tr}}(X) \rightarrow h(M)$. We define \mathbf{Rec} to be the full subcategory of \mathbf{PST} consisting of all objects having reciprocity.

3.2. Statement of the result and consequences.

Theorem 3.2.1. *Let $M = (\overline{X}, Y) \in \mathbf{MCor}$ be such that $X := \overline{X} \setminus |Y|$ is quasi-affine. Then $h_0(M) = h(M)$. Hence we have $\mathbf{RSC} \subset \mathbf{Rec}$.*

The proof of Theorem 3.2.1 will occupy §§3.3 and 3.4. We first deduce some consequences.

Corollary 3.2.2. *For any $F \in \mathbf{RSC}$, we have $F_{\operatorname{Zar}} \simeq F_{\operatorname{Nis}}$, where F_{Zar} (resp. F_{Nis}) is the Zariski (resp. Nisnevich) sheafification of F .*

Proof. Combine Theorem 3.2.1 and [7, Theorem 7]. \square

The next result depends on Theorem 2.4.1 (1).

Corollary 3.2.3. *Assume k is perfect. Then we have $\mathbf{RSC}_{\operatorname{Nis}} = \mathbf{Rec}_{\operatorname{Nis}}$.*

Proof. The inclusion follows immediately from Theorem 3.2.1. To prove the equality, let $F \in \mathbf{Rec}_{\operatorname{Nis}}$. By (2.5) and Theorem 3.2.1, the map $i^\sharp \rho F \rightarrow F$ of (2.6) is an isomorphism when evaluated at X if X is quasi-affine. By Theorem 2.4.3 (1), this extends to any $X \in \mathbf{Sm}$ by using a quasi-affine Zariski cover. Thus $F \in \mathbf{RSC}_{\operatorname{Nis}}$. \square

Remark 3.2.4. Here is an example of an object $F \in \mathbf{Rec} \setminus \mathbf{RSC}$. Define F as

$$\operatorname{Coker} \left(\bigoplus_{(X, a)} \mathbb{Z}_{\operatorname{tr}}(X) \rightarrow \mathbb{Z}_{\operatorname{tr}}(\mathbf{P}^1) \right)$$

where X runs through all smooth quasi-affine k -schemes and a runs through all elements of $\mathbf{Cor}(X, \mathbf{P}^1)$. By construction, $F(X) = 0$ for any smooth quasi-affine X , hence $F \in \mathbf{Rec}$. On the other hand, we claim that the image $\eta \in F(\mathbf{P}^1)$ of the identity map $1_{\mathbf{P}^1} \in \mathbb{Z}_{\operatorname{tr}}(\mathbf{P}^1)(\mathbf{P}^1)$

does not have an SC modulus. Since \mathbf{P}^1 is proper, this amounts to say that the composition

$$\underline{\mathrm{Hom}}(\mathbb{Z}_{\mathrm{tr}}(\mathbf{A}^1), \mathbb{Z}_{\mathrm{tr}}(\mathbf{P}^1)) \xrightarrow{i_0^* - i_1^*} \mathbb{Z}_{\mathrm{tr}}(\mathbf{P}^1) \xrightarrow{\eta} F$$

is nonzero. The quasi-affineness of the X 's yields that for any proper $Y \in \mathbf{Sm}$ the image of $\bigoplus_{(X,a)} \mathbb{Z}_{\mathrm{tr}}(X)(Y) \rightarrow \mathbb{Z}_{\mathrm{tr}}(\mathbf{P}^1)(Y) = \mathbf{Cor}(Y, \mathbf{P}^1)$ is generated by cycles of the form $Y \times x$ where x ranges over closed points of \mathbf{P}^1 . In particular, if we take $Y = \mathbf{P}^1$ we find that $F(\mathbf{P}^1)$ is not finitely generated. On the other hand, [10, Th. 3.3.1] shows

$$\begin{aligned} \mathrm{Coker}(\underline{\mathrm{Hom}}(\mathbb{Z}_{\mathrm{tr}}(\mathbf{A}^1), \mathbb{Z}_{\mathrm{tr}}(\mathbf{P}^1))(\mathbf{P}^1) \xrightarrow{i_0^* - i_1^*} \mathbb{Z}_{\mathrm{tr}}(\mathbf{P}^1)(\mathbf{P}^1)) \\ \simeq \mathrm{Pic}(\mathbf{P}^1 \times \mathbf{P}^1) \simeq \mathbb{Z} \times \mathbb{Z}. \end{aligned}$$

Hence η cannot vanish at \mathbf{P}^1 .

Corollary 3.2.5. *Assume k is perfect.*

- (1) *A presheaf with transfers represented by a smooth commutative algebraic group has SC-reciprocity.*
- (2) *The presheaf with transfers $H^0(-, \Omega_-^i)$ has SC-reciprocity for any $i \geq 0$. The same is true for the presheaf with transfers $H^0(-, \Omega_{-/k}^i)$.*
- (3) *Suppose that k is of positive characteristic. Then the presheaf with transfers $H^0(-, W_n \Omega_-^i)$ has SC-reciprocity for any $i \geq 0$ and $n \geq 1$.*

Proof. Combine Corollary 3.2.3 and [7, Theorems 4, 5]. \square

The next corollary uses the work of Binda et al [1]: we suppose k is of characteristic $p > 0$ and we use the notation $[1/p]$ to designate categories constructed out of sheaves of $\mathbb{Z}[1/p]$ -modules: they are full subcategories of those considered in this paper.

Corollary 3.2.6. *Assume that $\mathrm{char} k = p > 0$. Then the functor h_0^{rec} from Proposition 2.3.9 induces an equivalence of categories*

$$\mathbf{RSC}_{\mathrm{Nis}}[1/p] \xrightarrow{\sim} \mathbf{HI}_{\mathrm{Nis}}[1/p].$$

Proof. By Proposition 2.3.9, it suffices to show that $F \xrightarrow{\sim} h_0^{\mathrm{rec}}(F)$ for any $F \in \mathbf{RSC}_{\mathrm{Nis}}[1/p]$. If $F \in \mathbf{Rec}_{\mathrm{Nis}}[1/p]$, this follows from [1, Th. 3.5 (2)], hence the claim when k is perfect by Corollary 3.2.3. The general case reduces to this one by [3, Prop. 4.5]. \square

Remarks 3.2.7. There is a finer operation which consists of inverting p on morphisms rather than on objects, but Corollary 3.2.6 is false for these categories. For example, the sheaf $\bigoplus_{n \geq 1} W_n$ is a non-zero object of $\mathbf{RSC}_{\mathrm{Nis}}$, but h_0^{rec} maps it to 0 in $\mathbf{HI}_{\mathrm{Nis}}$.

In the sequel, (\bar{X}, Y) is as in Theorem 3.2.1.

3.3. Preliminary lemmas. In the rest of this section, we use a change of coordinates $\square \simeq (\mathbf{P}^1, 1)$ given by $\mathbf{A}^1 \rightarrow \mathbf{P}^1 \setminus \{1\}$, $t \mapsto t/(t-1)$. Let $\square := \mathbf{P}^1 - \{1\}$. Take $S \in \mathbf{Sm}$ and a closed integral subscheme $V \subset S \times \square \times X$ that is finite and surjective over $S \times \square$. We have a commutative diagram

$$(3.2) \quad \begin{array}{ccccc} & & V \hookrightarrow & X \times \square \times S & \\ & & \downarrow & & \downarrow \\ \bar{V}^N & \longrightarrow & \bar{V} \hookrightarrow & \bar{X} \times \mathbf{P}^1 \times S & \\ \downarrow & & \downarrow & & \downarrow p \\ \bar{W}^N & \longrightarrow & \bar{W} \hookrightarrow & \bar{X} \times S & \\ & & \searrow \gamma & & \downarrow \\ & & & & \bar{X} \end{array}$$

where \bar{V} is the closure of V in $\bar{X} \times \mathbf{P}^1 \times S$, \bar{W} is the image of \bar{V} under the projection p , and $\bar{V}^N \rightarrow \bar{V}$ and $\bar{W}^N \rightarrow \bar{W}$ are the normalizations. Let $\varphi_V : \bar{V}^N \rightarrow \bar{X} \times \mathbf{P}^1 \times S$ be the natural map. Let $\iota_\infty : \bar{X} \times S \rightarrow \bar{X} \times \mathbf{P}^1 \times S$ be induced by $\infty \in \mathbf{P}^1$. Put

$$\partial^\infty \bar{V} = \iota_\infty^{-1}(\bar{V}) = p(\bar{V} \cap (\bar{X} \times \{\infty\} \times S)) \subset \bar{X} \times S.$$

Putting $\bar{W}^o = \bar{W} \setminus \partial^\infty \bar{V}$ and $\bar{W}^{N,o} = \bar{W}^N \times_{\bar{W}} \bar{W}^o$, we have

$$(3.3) \quad \bar{V} \times_{\bar{W}} \bar{W}^{N,o} \subset \bar{W}^{N,o} \times (\mathbf{P}^1 - \{\infty\}).$$

Let \bar{V}^o be the reduced part of an irreducible component of $\bar{V} \times_{\bar{W}} \bar{W}^{N,o}$ which dominates $\bar{W}^{N,o}$. (Thus $\bar{V}^o \rightarrow \bar{V}$ is birational.)

Lemma 3.3.1. *If $\bar{W}^o = \emptyset$, then $V = W \times \square$ with $W = \bar{W} \cap (X \times S)$.*

Proof. The assumption implies $\bar{W} \subset p(\bar{V} \cap (\bar{X} \times \{\infty\} \times S))$ and hence

$$\dim \bar{W} \leq \dim \bar{V} \cap (\bar{X} \times \{\infty\} \times S) < \dim \bar{V}.$$

Noting $\bar{V} \hookrightarrow \bar{W} \times \mathbf{P}^1$, we get $\bar{V} = \bar{W} \times \mathbf{P}^1$, which implies the desired assertion. \square

Lemma 3.3.2. *If $\bar{W}^o \neq \emptyset$, \bar{V}^o is finite over $\bar{W}^{N,o}$.*

Proof. \bar{V} is proper over \bar{W} so that \bar{V}^o is proper over $\bar{W}^{N,o}$. On the other hand $\bar{W}^{N,o} \times (\mathbf{P}^1 - \{\infty\})$ is affine over $\bar{W}^{N,o}$ and so is \bar{V}^o . This implies the lemma. \square

Now we consider the modulus condition for V :

$$(3.4) \quad \varphi_V^{-1}(Y \times \mathbf{P}^1 \times S) \leq \varphi_V^{-1}(\overline{X} \times \{1\} \times S).$$

Let y be the standard coordinate on $\mathbf{P}^1 - \{\infty\} = \text{Spec}(k[y])$. (Note that the divisor involved in the modulus condition is $\{1\} \subset \mathbf{P}^1 - \{\infty\}$ defined by the ideal $(1 - y) \subset k[y]$.) Let $I \subset \mathcal{O}_{\overline{W}^{N,o}}$ be the ideal sheaf of $Y \times_{\overline{X}} \overline{W}^{N,o} \subset \overline{W}^{N,o}$.

Lemma 3.3.3. *Assuming $\overline{W}^o \neq \emptyset$, (3.4) is equivalent to the conditions:*

(i) $\overline{V} \cap (Y \times \square \times S) = \emptyset$.

(ii) *Locally on $\overline{W}^{N,o}$, \overline{V}^o is defined by an equation*

$$f(y) := (1 - y)^m + \sum_{1 \leq \nu \leq m} a_\nu (1 - y)^{m-\nu} \quad \text{with } a_\nu \in \Gamma(\overline{W}^{N,o}, I^\nu),$$

$$\text{in } \overline{W}^{N,o} \times (\mathbf{P}^1 - \{1\}) = \overline{W}^{N,o} \times \text{Spec}(k[y]) \text{ (see (3.3)).}$$

Proof. By Lemma 3.3.2, the minimal polynomial over $k(\overline{W})$ of the image of y in $\Gamma(\overline{V}^o, \mathcal{O})$:

$$f(t) = (1 - t)^m + \sum_{1 \leq \nu \leq m} a_\nu (1 - t)^{m-\nu}$$

has its coefficients $a_\nu \in A := \Gamma(\overline{W}^{N,o}, \mathcal{O})$. We claim that \overline{V}^o coincides with the closed subscheme $T \subset \overline{W}^{N,o} \times \text{Spec}(k[y])$ defined by the equation $f(y) \in A[y]$. Indeed it is clear that \overline{V}^o is contained in T , hence it suffices to show that T is integral. Note that T is a Cartier divisor in $\overline{W}^{N,o} \times (\mathbf{P}^1 - \{1\})$ which is finite over $\overline{W}^{N,o}$. It follows that each irreducible component dominates $\overline{W}^{N,o}$. Hence the integrality is checked over the generic point, which holds by the irreducibility of f . The claim is proved. Thus we are reduced to showing the following.

Claim 3.3.4. The condition (3.4) holds if and only if $\overline{V} \cap (Y \times \square \times S) = \emptyset$ and $a_\nu \in \Gamma(\overline{W}^{N,o}, I^\nu)$ for all ν .

The question is Zariski local and we may assume that I is generated by $\pi \in \Gamma(\overline{W}^{N,o}, \mathcal{O})$. Then (3.4) holds if and only if $\overline{V} \cap (Y \times \square \times S) = \emptyset$ and

$$(3.5) \quad \theta := \frac{1 - \bar{y}}{\pi} \in \Gamma(\overline{V}^N \times_{\overline{W}^N} \overline{W}^{N,o}, \mathcal{O}).$$

Noting $\pi \in k(\overline{W})$, the minimal polynomial of θ over $k(\overline{W})$ is

$$g(t) = t^m + \sum_{1 \leq \nu \leq m} \frac{a_\nu}{\pi^\nu} t^{m-\nu}.$$

Since \overline{V}^o is finite over $\overline{W}^{N,o}$ as is shown before, $\overline{V}^N \times_{\overline{W}^N} \overline{W}^{N,o}$ is finite over $\overline{W}^{N,o}$. Hence (3.5) is equivalent to the condition that θ is integral over $\Gamma(\overline{W}^{N,o}, \mathcal{O})$, which is equivalent to

$$\frac{a_\nu}{\pi^\nu} \in \Gamma(\overline{W}^{N,o}, \mathcal{O}) \quad \text{for all } \nu.$$

This proves the claim and the proof of Lemma 3.3.3 is completed. \square

3.4. Proof of Theorem 3.2.1. We put

$$C_1(\overline{X}|Y) := \omega_! \underline{\mathrm{Hom}}_{\mathbf{MPST}}(\mathbb{Z}_{\mathrm{tr}}(\overline{\square}), \mathbb{Z}_{\mathrm{tr}}(M)) \in \mathbf{PST}$$

and write by ∂ for the boundary map $\delta_{1,0}^{0*} - \delta_{1,\infty}^{0*} : C_1(\overline{X}|Y) \rightarrow \mathbb{Z}_{\mathrm{tr}}(M)$. Fix $S \in \mathbf{Sm}$. By Definitions 2.2.4 and 3.1.1, it suffices to construct a homomorphism :

$$(3.6) \quad \xi : C_1(\overline{X}|Y)(S) \rightarrow \Phi(\overline{X}, Y)(S)$$

such that the following diagram commutes:

$$(3.7) \quad \begin{array}{ccc} C_1(\overline{X}|Y)(S) & \xrightarrow{\partial} & \mathbb{Z}_{\mathrm{tr}}(X)(S) \\ \downarrow \xi & & \parallel \\ \Phi(\overline{X}, Y)(S) & \xrightarrow{\tau} & \mathbb{Z}_{\mathrm{tr}}(X)(S) \end{array}$$

and such that we have

$$(3.8) \quad \mathrm{Image}(\tau) = \mathrm{Image}(\tau \circ \xi).$$

Take a closed integral subscheme $V \subset S \times \square \times X$, finite and surjective over $S \times \square$ and satisfying (3.4). Consider the commutative diagram (3.2) and let $\varphi : \overline{W}^N \rightarrow \overline{X} \times S$ be the induced map. We first suppose $\overline{W}^o \neq \emptyset$. Then we have (see §3.1 for notations)

$$(\overline{W}^N \xrightarrow{\varphi} \overline{X} \times S) \in \mathcal{C}_{(\overline{X}, Y)}(S).$$

The projection $V \rightarrow \square = \mathbf{P}^1 - \{1\}$ induces a rational function $g_V \in k(\overline{V})^\times$. By [2, Prop.1.4 and §1.6] we have

$$(3.9) \quad \partial V = \varphi_* \mathrm{div}_{\overline{W}^N}(Ng_V) \in \mathbb{Z}_{\mathrm{tr}}(X)(S) = \mathbf{Cor}(S, X),$$

where $N : k(\overline{V})^\times \rightarrow k(\overline{W})^\times$ is the norm map induced by $\overline{V} \rightarrow \overline{W}$. By Lemma 3.3.3 we have

$$Ng_V = f(0) = 1 + \sum_{1 \leq \nu \leq m} a_\nu \in \Gamma(\overline{W}^{N,o}, I) \subset G(\overline{W}^N, \gamma_\varphi^* Y) \subset \Phi(\overline{X}, Y)(S).$$

We now define a map

$$(3.10) \quad \xi : C_1(\overline{X}|Y)(S) \rightarrow \Phi(\overline{X}, Y)(S)$$

by declaring

$$\xi(V) = \begin{cases} Ng_V & \text{if } \overline{W}^o \neq \emptyset; \\ 0 & \text{if } \overline{W}^o = \emptyset. \end{cases}$$

Note that if $\overline{W}^o = \emptyset$, then we have $\partial(V) = 0$ by Lemma 3.3.1. It follows that the diagram (3.7) commutes thanks to (3.9).

It remains to show (3.8). To this end, we take $(\varphi_0 : \overline{C} \rightarrow \overline{X} \times S) \in \mathcal{C}_{(\overline{X}, Y)}(S)$ and show $\tau(G(\overline{C}, \gamma_{\varphi_0}^* Y)) \subset \text{Image}(\tau \circ \xi)$ (see §3.1 for notations). Let $\overline{W} \hookrightarrow \overline{X} \times S$ be the image of φ_0 and let $\overline{W}^N \rightarrow \overline{W}$ be its normalization so that $(\varphi : \overline{W}^N \rightarrow \overline{X} \times S) \in \mathcal{C}_{(\overline{X}, Y)}(S)$. Since $\tau(G(\overline{C}, \gamma_{\varphi_0}^* Y)) \subset \tau(G(\overline{W}^N, \gamma_{\varphi}^* Y))$, it suffices to show the following.

Lemma 3.4.1. *The subgroup $G(\overline{W}^N, \gamma_{\varphi}^* Y) \subset \Phi(\overline{X}, Y)(S)$ is contained in the image of $\xi : C_1(\overline{X}|Y)(S) \rightarrow \Phi(\overline{X}, Y)(S)$.*

Proof. Take $g \in G(\overline{W}^N, \gamma_{\varphi}^* Y)$. Let $\Sigma \subset \overline{W}^N$ be the closure of the union of points $x \in \overline{W}^N$ of codimension one such that $v_x(g) < 0$, where v_x is the valuation associated to x . Since \overline{W}^N is normal, we have $g \in \Gamma(\overline{W}^N - \Sigma, \mathcal{O})$ and $g \in G(\overline{W}^N, \gamma_{\varphi}^* Y)$ implies

$$(3.11) \quad g - 1 \in \Gamma(\overline{W}^N - \Sigma, I),$$

where $I \subset \mathcal{O}_{\overline{W}^N}$ is the ideal sheaf of $\gamma_{\varphi}^* Y \subset \overline{W}^N$. Let

$$\psi_g : \overline{W}^N - \Sigma \rightarrow \mathbf{P}^1 - \{\infty\}$$

be the morphism induced by g and $\Gamma \subset \overline{W}^N \times \mathbf{P}^1$ be the closure of the graph of ψ_g . Let

$$\overline{V} \subset \overline{W} \times \mathbf{P}^1 \subset \overline{X} \times \mathbf{P}^1 \times S$$

be the image of Γ under $\overline{W}^N \times \mathbf{P}^1 \rightarrow \overline{W} \times \mathbf{P}^1$. By (3.11) we have $|\gamma_{\varphi}^* Y| \subset \psi_g^{-1}(1)$ and hence

$$(3.12) \quad \overline{V} \cap (Y \times \square \times S) = \emptyset$$

so that

$$V := \overline{V} \cap (\overline{X} \times \square \times S) \subset X \times \square \times S.$$

It suffices to show the following.

Claim 3.4.2. $V \in C_1(\overline{X}|Y)(S)$ and $\xi(V) = g$.

Once we prove the first assertion, the second follows easily from the construction of ξ . To prove the first assertion, by (3.12), the map $V \rightarrow \square \times S$ is proper and hence finite since X is quasi-affine by the assumption. Moreover it is surjective since $\dim V = \dim \overline{W} = \dim S +$

1. Hence it suffices to check the condition (ii) of Lemma 3.3.3. By definition

(♠) $\Gamma \cap ((\overline{W}^N - \Sigma) \times (\mathbf{P}^1 - \{\infty\}))$ is the graph of ψ_g and hence is defined by $y - g$ where y is the standard coordinate of $\mathbf{P}^1 - \{\infty\} = \text{Spec}(k[y])$.

We have a diagram of schemes

$$(3.13) \quad \begin{array}{ccccc} \overline{W} & \xrightarrow{\iota_\infty} & \overline{W} \times \mathbf{P}^1 & \longleftarrow & \overline{V} \\ \pi_{\overline{W}} \uparrow & & \uparrow & & \uparrow \pi_{\overline{V}} \\ \overline{W}^N & \xrightarrow{\iota_\infty} & \overline{W}^N \times \mathbf{P}^1 & \longleftarrow & \Gamma' := \overline{V} \times_{\overline{W}} \overline{W}^N \\ & \searrow id_{\overline{W}} & \downarrow pr & \swarrow pr_\Gamma & \swarrow \\ & & \overline{W}^N & & \end{array}$$

where ι_∞ are induced by $\infty \in \mathbf{P}^1$. The natural map $\Gamma \rightarrow \Gamma'$ is a closed immersion onto an irreducible component that dominates \overline{V} . We claim

$$(3.14) \quad \Sigma \subset \iota_\infty^{-1}(\Gamma).$$

The claim implies $\overline{W}^{N,o} := \overline{W}^N \times_{\overline{W}} (\overline{W} \setminus \iota_\infty^{-1}(\overline{V})) \subset \overline{W}^N - \Sigma$. Let \overline{V}^o be the reduced part of the irreducible component of

$$\overline{V}^{o'} := \overline{V} \times_{\overline{W}} \overline{W}^{N,o} = \Gamma' \times_{\overline{W}^N} \overline{W}^{N,o} \subset \overline{W}^{N,o} \times (\mathbf{P}^1 - \{\infty\})$$

which dominates \overline{V} ; see the following diagram:

$$\begin{array}{ccccccc} & & \overline{W}^{N,o} & \longleftarrow & \overline{W}^{N,o} \times (\mathbf{P}^1 - \{\infty\}) & \longleftarrow & \overline{V}^{o'} & \longleftarrow & \overline{V}^o \\ & \swarrow & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \overline{W}^N - \Sigma & \hookrightarrow & \overline{W}^N & \longleftarrow & \overline{W}^N \times \mathbf{P}^1 & \longleftarrow & \Gamma' & \longleftarrow & \Gamma \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ & & \overline{W} & \longleftarrow & \overline{W} \times \mathbf{P}^1 & \longleftarrow & \overline{V} & & \end{array}$$

By (♠), \overline{V}^o is defined in $\overline{W}^{N,o} \times \text{Spec}(k[y])$ by the equation $y - g$ and thus V satisfies Lemma 3.3.3 (ii).

It remains to show (3.14). From (3.13), it is equivalent to

$$(3.15) \quad \Sigma \subset pr((\overline{W}^N \times \infty) \cap \Gamma).$$

Since pr_Γ is proper birational and \overline{W}^N is normal, pr_Γ is an isomorphism above all codimension one points in \overline{W}^N (but not necessarily in all codimension one points of Γ). For a generic point $x \in \Sigma$, there is a unique codimension one point $y \in \Gamma$ such that $x = pr_\Gamma(y)$ and we have

$v_y(g) = v_x(g) < 0$ for $g \in k(\Gamma) = k(\overline{W}^N)$. The projection $\overline{W}^N \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ induces a morphism $\Gamma \setminus (\overline{W}^N \times \{\infty\}) \rightarrow \mathbf{P}^1 - \{\infty\}$, which corresponds to g . Hence we must have $y \in (\overline{W}^N \times \{\infty\}) \cap \Gamma$ which proves (3.15) by the properness of pr_Γ . This completes the proof of Lemma 3.4.1. \square

REFERENCES

- [1] F. Binda, J. Cao, W. Kai, R. Sugiyama, *Torsion and divisibility for reciprocity sheaves and 0-cycles with modulus*, J. Alg. **469** (2017), 437–463.
- [2] W. Fulton, *Intersection Theory*, Ergebnisse der Math. und ihrer Grenzgebiete 3. Folge **2**, Springer-Verlag, 1984.
- [3] B. Kahn, *Motifs et adjoints*, Rendic. Sem. Math. Univ. Padova **139** (2018), 77–128.
- [4] B. Kahn, H. Miyazaki, S. Saito, T. Yamazaki, *Motives with modulus, I: Modulus sheaves with transfers for non-proper modulus pairs*, <https://arxiv.org/abs/1908.02975>, Épijournal de Géométrie Algébrique **5** (2021), Art. 1, 46 pp.
- [5] B. Kahn, H. Miyazaki, S. Saito, T. Yamazaki, *Motives with modulus, II: Modulus sheaves with transfers for proper modulus pairs*, <https://arxiv.org/abs/1910.14534>, Épijournal de Géométrie Algébrique **5** (2021), Art. 2, 31 pp.
- [6] B. Kahn, H. Miyazaki, S. Saito, T. Yamazaki, *Motives with modulus, III: The categories of motives*, preprint, 2020, <https://arxiv.org/abs/2011.11859>.
- [7] B. Kahn, S. Saito, T. Yamazaki, *Reciprocity sheaves* (with two appendices by Kay Rülling), Compos. Math. **152** (2016), no. 9, 1851–1898.
- [8] B. Kahn, S. Saito, T. Yamazaki, *Motives with modulus*, <https://arxiv.org/abs/1511.07124>, withdrawn.
- [9] B. Kahn, S. Saito, T. Yamazaki, *Reciprocity sheaves and motives with modulus*, <https://arxiv.org/pdf/1707.07398v3.pdf>.
- [10] B. Kahn, R. Sujatha, *Birational motives, II: triangulated birational motives*, IMRN **2017**, 6778–6831.
- [11] B. Kahn, T. Yamazaki, *Voevodsky’s motives and Weil reciprocity*, Duke Math. J. **62** (2013), 2751–2796.
- [12] M. Kerz, S. Saito, *Chow group of 0-cycles with modulus and higher dimensional class field theory*, Duke Math. J. **165**, no. 15 (2016), 2811–28974.
- [13] C. Mazza, V. Voevodsky, C. Weibel, *Lecture notes on motivic cohomology*, Clay Math. Monographs **2**, AMS, 2006.
- [14] K. Rülling and S. Saito, *Reciprocity sheaves and abelian ramification theory*, preprint, 2018, <https://arxiv.org/abs/1812.08716>.
- [15] K. Rülling, T. Yamazaki, *Suslin homology of relative curves with modulus*, J. Lond. Math. Soc. (2) **93** (2016), no. 3, 567–589.
- [16] S. Saito, *Purity of reciprocity sheaves*, Adv. Math. **366** (2020), 107067, 70 pp.
- [17] J.-P. Serre, *Groupes algébriques et corps de classes*, Hermann, 1959.
- [18] V. Voevodsky, *Triangulated categories of motives over a field*, in E. Friedlander, A. Suslin, V. Voevodsky Cycles, transfers and motivic cohomology theories, Ann. Math. Studies **143**, Princeton University Press, 2000, 188–238.

IMJ-PRG, CASE 247, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
Email address: `bruno.kahn@imj-prg.fr`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO,
3-8-1 KOMABA,, TOKYO 153-8941, JAPAN
Email address: `sshuji@msb.biglobe.ne.jp`

INSTITUTE OF MATHEMATICS, TOHOKU UNIVERSITY, AOBA, SENDAI 980-
8578, JAPAN
Email address: `takao.yamazaki.b6@tohoku.ac.jp`