

CYCLE CLASS MAPS FOR CHOW GROUPS OF ZERO-CYCLES WITH MODULUS

KAY RÜLLING AND SHUJI SAITO

ABSTRACT. For a quasi-projective smooth scheme X of pure dimension d over a field k and an effective Cartier divisor $D \subset X$ whose support is a simple normal crossing divisor, we construct a cycle class map

$$cyc_{X|D} : \mathrm{CH}_0(X|D) \rightarrow H_{\mathrm{Nis}}^d(X, K_d^M(\mathcal{O}_X, I_D))$$

from the Chow group of zero-cycles with modulus to the cohomology of the relative Milnor K -sheaf.

CONTENTS

1. Introduction	1
2. The projective bundle formula for relative Milnor K -theory	3
3. The cycle class map	12
References	14

1. INTRODUCTION

Let U be a smooth variety of dimension d over a finite field k and $U \hookrightarrow X$ be an open immersion with X normal and proper over k such that $X \setminus U$ is the support of an effective Cartier divisor on X . Let $\pi_1^{\mathrm{ab}}(U)$ be the abelianized fundamental group of U . Kato and Saito [KS86] established class field theory for U by using the reciprocity map

$$\rho_U : \varprojlim_D H^d(X_{\mathrm{Nis}}, K_d^M(\mathcal{O}_X, I_D)) \rightarrow \pi_1^{\mathrm{ab}}(U),$$

where the limits are over all effective Cartier divisors D supported on $X \setminus U$, and $K_d^M(\mathcal{O}_X, I_D)$ with $I_D \subset \mathcal{O}_X$ the ideal sheaf of D is the relative Milnor K -sheaf on the Nisnevich site over X (see (2.15.2)). A refinement of this statement was given by Kerz-Saito [KS16]¹ using another reciprocity map

$$\phi_U : \varprojlim_D \mathrm{CH}_0(X|D) \rightarrow \pi_1^{\mathrm{ab}}(U),$$

where $\mathrm{CH}_0(X|D)$ is the Chow group of zero-cycles with modulus. By definition $\mathrm{CH}_0(X|D)$ is the quotient of the group $Z_0(U)$ of zero-cycles on U by the subgroup generated by divisors of rational functions on curves in U which satisfy a certain modulus conditions with respect to D (see 3.1). On the other hand, the Gersten resolution of the Milnor K -sheaf yields a natural map

$$(1.0.1) \quad \theta : Z_0(U) \longrightarrow H^d(X_{\mathrm{Nis}}, K_d^M(\mathcal{O}_X, I_D)),$$

K.R. was supported by the DFG Heisenberg Grant RU 1412/2-2. S.S. is supported by the JSPS KAKENHI Grant (20H01791).

¹They needed to assume $\mathrm{ch}(k) \neq 2$. This assumption was removed by Binda-Krishna-Saito [BKS21] giving a simpler proof of the main result in [KS16].

which is surjective by [KS86], see (3.1.3).

With the help of ramification theory, the main result of [KS16] implies that ϕ_U induces a map for a fixed divisor D :

$$\phi_{X|D} : \mathrm{CH}_0(X|D) \rightarrow \pi_1^{\mathrm{ab}}(X|D)$$

whose restriction to the degree-zero part is an isomorphism of finite abelian groups. Here, $\pi_1^{\mathrm{ab}}(X|D)$ is the quotient of $\pi_1^{\mathrm{ab}}(U)$ which classifies abelian étale coverings of U with ramification over $X \setminus U$ bounded by D , see [KS14]. A natural question is whether the latter fact holds with $\mathrm{CH}_0(X|D)$ replaced by $H^d(X_{\mathrm{Nis}}, K_d^M(\mathcal{O}_X, I_D))$. A positive answer was given by Gupta-Krishna [GK21] assuming X is smooth and projective over the finite field k . It implies that there is a natural isomorphism

$$\mathrm{cyc}_{X|D} : \mathrm{CH}_0(X|D) \cong H_{\mathrm{Nis}}^d(X, K_d^M(\mathcal{O}_X, I_D)),$$

which gives a factorization of θ from (1.0.1), namely fits into the following commutative diagram

$$(1.0.2) \quad \begin{array}{ccc} & Z_0(U) & \\ \pi \swarrow & & \searrow \theta \\ \mathrm{CH}_0(X|D) & \xrightarrow{\mathrm{cyc}_{X|D}} & H_{\mathrm{Nis}}^d(X, K_d^M(\mathcal{O}_X, I_D)) \end{array}$$

where π is the quotient map. The question whether such a factorization of θ via $\mathrm{CH}_0(X|D)$ exists makes sense in case X is smooth (not necessarily projective) over an arbitrary field k and D is an effective Cartier divisor on X . It was positively answered in the following cases in which X is always smooth and D is an effective Cartier divisor:

- (1) X is a quasi-projective surface over a field, see [Kri15, Theorem 1.2];
- (2) X is affine over an algebraically closed field, see [GK20, Theorem 1.3(1)];
- (3) X is projective over an algebraically closed field and D is integral, see [GK20, Theorem 1.3(2)];

In all the above cases the cycle map $\mathrm{cyc}_{X|D}$ is actually an isomorphism, by [BKS21] in the case (1) and by [GK20] in the cases (1), (2). The authors were furthermore informed by Krishna that the existence of the cycle map also follows for smooth projective varieties over algebraically closed fields if D is reduced from [Kri18].

In this note we prove:

Theorem 1.1 (Theorem 3.2). *Let X be a quasi-projective smooth scheme of pure dimension d over a field k and $D \subset X$ be an effective Cartier divisor whose support is a simple normal crossing divisor. There exists a map $\mathrm{cyc}_{X|D}$ which makes (1.0.2) commutative.*

We will use the cycle map from the above theorem in [RS21a], where it is important to allow an arbitrary base field. This is our main motivation.

The proof of Theorem 1.1 is different in spirit from the proofs in (1)-(3) above and follows instead the same strategy as in [RS18], in which a factorization

$$(1.1.1) \quad \mathrm{CH}_0(X|D) \rightarrow H_{\mathrm{Nis}}^d(X, U_{d,X|D})$$

is constructed, where $U_{d,X|D}$ is a variant of $K_d^M(\mathcal{O}_X, I_D)$ (see 2.1). There is another variant $V_{d,X|D}$ and it is easy to prove that $K_d^M(\mathcal{O}_X, I_D)$ and $V_{d,X|D}$ have the

same cohomology group in the top degree d (see (2.16.2)). Thus we are reduced to constructing a factorization

$$(1.1.2) \quad \mathrm{CH}_0(X|D) \rightarrow H_{\mathrm{Nis}}^d(X, V_{d,X|D}).$$

A key input in the construction of (1.1.1) is a projective bundle formula for the cohomology of $U_{d,X|D}$. In §2, we prove its variant for $V_{d,X|D}$. This is deduced from the original statement by comparing the cohomologies of $U_{d,X|D}$ and $V_{d,X|D}$. To this end we define a decreasing filtration on $V_{d,X|D}$, which ends in $U_{d,X|D+D_{\mathrm{red}}}$ (see 2.5) and compute the associated graded algebra, see Proposition 2.9. Having this refined projective bundle formula at hand, we construct (1.1.2) in §3 using the same argument as for the construction of (1.1.1) in [RS18].

Acknowledgments. The authors thank Amalendu Krishna for his comments, in particular for pointing out the references [Kri15], [Kri18], and [GK20].

2. THE PROJECTIVE BUNDLE FORMULA FOR RELATIVE MILNOR K -THEORY

In this section k is a field of characteristic $p \geq 0$ and \mathbf{Sm} denotes the category of smooth k -schemes.

2.1. Let X be a k -scheme. We denote by $K_{r,X}^M$ ($r \geq 0$) the Nisnevich sheafification of the improved Milnor K -theory from [Ker10]. Let $D \subset X$ be a closed subscheme and denote by $j : U := X \setminus D \hookrightarrow X$ (resp. $i : D \hookrightarrow X$) the corresponding open (resp. closed) immersion. Set

$$\mathcal{O}_{X|D}^\times := \mathrm{Ker}(\mathcal{O}_X^\times \rightarrow i_*\mathcal{O}_D^\times).$$

We consider the following Nisnevich sheaves for $r \geq 1$

$$U_{r,X|D} := \mathrm{Im}(\mathcal{O}_{X|D}^\times \otimes_{\mathbb{Z}} j_* K_{r-1,U}^M \rightarrow j_* K_{r,U}^M), \quad V_{r,X|D} := \mathrm{Im}(\mathcal{O}_{X|D}^\times \otimes_{\mathbb{Z}} K_{r-1,X}^M \rightarrow K_{r,X}^M).$$

Note that

- (1) $U_{1,X|D} = V_{1,X|D} = \mathcal{O}_{X|D}^\times$ and there is a natural map $V_{r,X|D} \rightarrow U_{r,X|D}$, which is injective if X is regular (by the Gersten resolution, see [Ker10, Proposition 10(8)]);
- (2) if k is an infinite field, then $V_{r,X|D}$ is equal to the Nisnevich sheaf $\mathrm{Ker}(K_{r,X}^M \rightarrow K_{r,D}^M)$ considered in [KS86, (1.3)] (this follows from [KS86, Lemma 1.3.1] and [Ker10, Proposition 10(5)]);
- (3) the sheaf $U_{r,X|D}$ (for D an effective Cartier divisor) is equal to the Nisnevich sheaf $\mathcal{K}_{r,X|D,\mathrm{Nis}}^M$ considered in [RS18, 2.3].

2.2. Let $X \in \mathbf{Sm}$ and let $D = \mathrm{div}(t)$ be a smooth integral and principal divisor on X . For $m \geq 1$ and $r \geq 1$ we set

$$U_{r,m} = U_{r,X|mD}, \quad V_{r,m} = V_{r,X|mD}.$$

If $p > 0$, write $m = p^s m'$ with $(p, m') = 1$ and $s \geq 0$; if $p = 0$ we set $s = 0$ and $m' := m$. Denote by Ω_D^i the sheaf of absolute differential i -forms and set $B_0 \Omega_D^i = 0$ and $Z_0 \Omega_D^i = \Omega_D^i$. For $s \geq 1$ (hence $p > 0$), the sheaves $B_s \Omega_D^i$ and $Z_s \Omega_D^i$ are recursively defined by

$$C^{-1} : Z_{s-1} \Omega_D^i \xrightarrow{\cong} \frac{Z_s \Omega_D^i}{d\Omega_D^{i-1}}, \quad C^{-1} : B_{s-1} \Omega_D^i \xrightarrow{\cong} \frac{B_s \Omega_D^i}{d\Omega_D^{i-1}},$$

where C^{-1} is inverse Cartier operator.

Set

$${}^m G^r := \text{Coker} \left(\Omega_D^{r-2} \rightarrow \frac{\Omega_D^{r-1}}{B_s} \oplus \frac{\Omega_D^{r-2}}{B_s}, \quad \alpha \mapsto (C^{-s}(d\alpha), (-1)^r m' C^{-s}\alpha) \right),$$

where C^{-s} is the s -fold iterated inverse Cartier operator, for $s \geq 1$, and $C^{-0} = \text{id}$. By [BK86, Remark (4.8)] (see also [RS18, Proposition 2.15, Theorem 2.19], in particular for $s = 0$) we have an isomorphism

$$(2.2.1) \quad {}^m G^r \xrightarrow{\cong} \frac{U_{r,m}}{U_{r,m+1}}$$

given by

$$(a \cdot \text{dlog } \underline{b}, e \cdot \text{dlog } \underline{f}) \mapsto \{1 + t^m \tilde{a}, \tilde{b}\} + \{1 + t^m \tilde{e}, \tilde{f}, t\},$$

where $a, e \in \mathcal{O}_D$, $\underline{b} = (b_1, \dots, b_{r-1})$, and $\underline{f} = (f_1, \dots, f_{r-2})$, $b_i, f_j \in \mathcal{O}_D^\times$ are local sections and $\tilde{a}, \tilde{e}, \tilde{b}_i, \tilde{f}_j$ are lifts to \mathcal{O}_X .

Lemma 2.3. *Let the assumption and notation be as in 2.2. There is an isomorphism of exact sequences of sheaves on D*

$$(2.3.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \frac{V_{r,m}}{U_{r,m+1}} & \longrightarrow & \frac{U_{r,m}}{U_{r,m+1}} & \longrightarrow & \frac{U_{r,m}}{V_{r,m}} \longrightarrow 0 \\ & & \uparrow \cong & & \uparrow \cong (2.2.1) & & \uparrow \cong \\ 0 & \longrightarrow & \frac{\Omega_D^{r-1}}{B_s \Omega_D^{r-1}} & \longrightarrow & {}^m G^r & \longrightarrow & \frac{\Omega_D^{r-2}}{Z_s \Omega_D^{r-2}} \longrightarrow 0. \end{array}$$

In particular in characteristic 0, we have $U_{r,m} = V_{r,m}$.

Proof. Note that by the definition of (2.2.1) the composition

$$\Omega_D^{r-1}/B_s \rightarrow {}^m G^r \rightarrow U_{r,m}/U_{r,m+1}$$

factors via a surjection

$$\Omega_D^{r-1}/B_s \twoheadrightarrow V_{r,m}/U_{r,m+1}.$$

Therefore it suffices to show that the lower sequence is exact. This is immediate for $s = 0$. We assume $s \geq 1$ (hence $p > 0$). By [Ill79, I, Proposition 3.11] we have

$$B_s \Omega_D^{r-1} = F^{s-1} dW_s \Omega_D^{r-1}, \quad Z_s \Omega_D^{r-1} = F^s W_{s+1} \Omega_D^{r-1},$$

where $W_n \Omega_D^\bullet$ denotes the de Rham-Witt complex from [Ill79]. The lower row in (2.3.1) becomes

$$\frac{\Omega_D^{r-1}}{\{F^{s-1} d\gamma\}} \xrightarrow{\varphi} \frac{\Omega_D^{r-1} \oplus \Omega_D^{r-2}}{\{(F^s d\alpha, (-1)^r m' F^s(\alpha) + F^s d\beta)\}} \xrightarrow{\psi} \frac{\Omega_D^{r-2}}{\{F^s(\delta)\}},$$

where,

$$\varphi(x) = (x, 0), \quad \psi(x, y) = y.$$

Note that φ is well-defined since $F^{s-1} d\gamma = F^s dV\gamma$ and it clearly surjects onto the kernel of ψ . We show that φ is injective. Indeed, $\varphi(x) = 0$ implies that there exist α, β with $x = F^s d\alpha$ and $F^s \alpha = F^s d\beta$; since $p^s = V^s F^s$ we obtain $p^s(\alpha - d\beta) = 0$; hence by [Ill79, I, Proposition 3.4] we find γ, δ such that $\alpha - d\beta = V(\gamma) + dV(\delta)$; thus

$$x = F^s d\alpha = F^s dV(\gamma) + F^s dd(\beta + V(\delta)) = F^{s-1} d\gamma.$$

Therefore the lower sequence is exact. \square

The following lemma is direct to check (see [RS18, Lemma 2.7]).

Lemma 2.4. *Let R be an integral local ring with maximal ideal \mathfrak{m} and fraction field K . Let $a, b, c \in R$ and $s, t \in \mathfrak{m}$. The following equalities hold in $K_2^M(K)$*

- (1) $\{1 + as, 1 + bt\} = -\{1 + \frac{ab}{1+as}st, -as(1 + bt)\};$
- (2) $\{1 + \frac{s-1}{1+ct}ct, 1 - \frac{1+ct}{1+cst}s\} = \{1 + cst, s\}$

2.5. Let $X \in \mathbf{Sm}$ and let $D = \sum_{j=1}^n m_j D_j$ be an effective divisor, such that $|D| = \sum_{j=1}^n D_j$ is a simple normal crossing divisor with smooth irreducible components D_j and $m_j \geq 1$. For $i \geq 0$ we set

$$E_i := D_{i+1} + \dots + D_n,$$

in particular $E_n = 0$, and denote by $j_i : U_i := X \setminus |E_i| \hookrightarrow X$ the open immersion (with the notation from 2.1 we have $j = j_0 : U = U_0 \hookrightarrow X$). For $r \geq 1$ we define recursively

$$V_r^{-1} := 0 \quad \text{and} \quad V_r^i := V_r^{i-1} + \text{Im} \left(\mathcal{O}_{X|D+E_i}^\times \otimes_{\mathbb{Z}} j_{i*} K_{r-1, X \setminus E_i}^M \rightarrow j_{i*} K_{r, U}^M \right), \quad i \geq 0.$$

By Lemma 2.4(2) we have $V_r^i \subset V_{r, X|D}$, for all i . We obtain a filtration

$$U_{r, X|D+E_0} = V_r^0 \subset V_r^1 \subset \dots \subset V_r^n = V_{r, X|D}.$$

2.6. Let $r \geq 1$. Let $X \in \mathbf{Sm}$ and let $D = \sum_{j=1}^n m_j D_j$ be as in 2.5. If $p > 0$, write $m_j = p^{s_j} m'_j$ with $(p, m'_j) = 1$ and $s_j \geq 0$; if $p = 0$ we set $m'_j := m_j$ and $s_j = 0$. (We will use the convention $0^0 = 1$.) After renumbering we can assume that

$$(2.6.1) \quad s_1 \geq s_2 \geq \dots \geq s_n \geq 0.$$

Fix $i \in \{1, \dots, n\}$. If $p > 0$, then there is an inverse Cartier operator on the absolute logarithmic differential forms

$$C^{-1} : \Omega_{D_i}^{r-1}(\log E_i) \rightarrow \frac{F_* \Omega_{D_i}^{r-1}(\log E_i)}{d\Omega_{D_i}^{r-2}(\log E_i)},$$

where F is the absolute Frobenius and we write $\Omega_{D_i}^{r-1}(\log E_i)$ short for $\Omega_{D_i}^{r-1}(\log E_i|_{D_i})$ etc. The \mathcal{O}_{D_i} -submodule $B_s \Omega_{D_i}^{r-1}(\log E_i) \subset F_* \Omega_{D_i}^{r-1}$ is recursively defined by

$$B_0 \Omega_{D_i}^{r-1}(\log E_i) := 0, \quad C^{-1} : B_{s-1} \Omega_{D_i}^{r-1}(\log E_i) \xrightarrow{\simeq} \frac{B_s \Omega_{D_i}^{r-1}(\log E_i)}{d\Omega_{D_i}^{r-2}(\log E_i)}.$$

Denote by $D(i)$ the minimal divisor satisfying $p^{s_i} D(i) \geq D + E_i$, i.e., by (2.6.1)

$$D(i) = \sum_{j=1}^i p^{s_j - s_i} m'_j D_j + \sum_{j=i+1}^n \left\lceil \frac{m_j + 1}{p^{s_i}} \right\rceil D_j.$$

We write $\mathcal{O}_{D_i}(-D) := \mathcal{O}_X(-D)|_{D_i}$ etc. and have natural inclusions of $\mathcal{O}_{D_i}^{p^{s_i}}$ -modules

$$\begin{aligned} \mathcal{O}_{D_i}(-D(i))^{p^{s_i}} \otimes_{\mathcal{O}_{D_i}^{p^{s_i}}} B_{s_i} \Omega_{D_i}^{r-1}(\log E_i) &\subset \mathcal{O}_{D_i}(-D(i))^{p^{s_i}} \otimes_{\mathcal{O}_{D_i}^{p^{s_i}}} \Omega_{D_i}^{r-1}(\log E_i) \\ &\subset \mathcal{O}_{D_i}(-D - E_i) \otimes_{\mathcal{O}_{D_i}} \Omega_{D_i}^{r-1}(\log E_i). \end{aligned}$$

We obtain a sheaf of $\mathcal{O}_{D_i}^{p^{s_i}}$ -modules

$$Q_D^{i, r-1} := \frac{\mathcal{O}_{D_i}(-D - E_i) \otimes_{\mathcal{O}_{D_i}} \Omega_{D_i}^{r-1}(\log E_i)}{\mathcal{O}_{D_i}(-D(i))^{p^{s_i}} \otimes_{\mathcal{O}_{D_i}^{p^{s_i}}} B_{s_i} \Omega_{D_i}^{r-1}(\log E_i)}.$$

If $p = 0$, then $s_i = 0$ and we have $Q_D^{i, r-1} = \mathcal{O}_{D_i}(-D - E_i) \otimes_{\mathcal{O}_{D_i}} \Omega_{D_i}^{r-1}(\log E_i)$.

Lemma 2.7. *Let the notations and assumptions be as in 2.6. Assume k is perfect.*

- (1) *Then $Q_D^{i,r-1}$ is a locally free \mathcal{O}_{D_i} -module.*
(2) *Let $j_i : U_i = D_i \setminus |E_i| \hookrightarrow D_i$ be the open immersion. Then*

$$j_i^* Q_D^{i,r-1} = \mathcal{O}_{U_i} \left(- \sum_{j=1}^i p^{s_j - s_i} m'_j D_j \right)^{p^{s_i}} \otimes_{\mathcal{O}_{U_i}^{p^{s_i}}} \frac{\Omega_{U_i}^{r-1}}{B_{s_i} \Omega_{U_i}^{r-1}}$$

and the restriction map $Q_D^{i,r-1} \rightarrow j_{i} j_i^* Q_D^{i,r-1}$ is injective.*

Proof. The first statement in (2) is clear, the second statement follows from (1). Furthermore, (1) is immediate if $s_i = 0$. Thus we assume $s_i \geq 1$ and hence also $p > 0$. The statement is local. Let $x \in X$ be a point, up to replacing $\{D_j\}_j$ by some ordered subset we can assume that all D_j contain x . Hence we may assume $X = \text{Spec } R$, with R a local ring, essentially smooth over k and with a regular sequence of parameters t_1, \dots, t_n , where t_j is a local equation for D_j , $j = 1, \dots, n$. Then $D_i = \text{Spec } R_i$, with $R_i = R/(t_i)$. Set

$$(2.7.1) \quad \tau := \prod_{j=1}^i t_j^{m_j} \prod_{j=i+1}^n t_j^{m_j+1},$$

which is an equation for $D + E_i$. Since k is perfect of positive characteristic, an $R_i^{p^{s_i}}$ -basis of $\tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$ is given by

$$\tau \cdot t_1^{j_1} \cdots t_N^{j_N} \cdot dt_{k_1} \cdots dt_{k_{r_1}} \cdot d\log t_{l_1} \cdots d\log t_{l_{r_2}},$$

where $j_\nu \in [0, p^{s_i} - 1]$, $1 \leq k_1 < \cdots < k_{r_1} \leq i$, $i+1 \leq l_1 < \cdots < l_{r_2} \leq n$, and $r_1, r_2 \geq 0$ with $r_1 + r_2 = r - 1$. By (2.6.1) we have a local equation for $p^{s_i} D(i)$ of the form

$$(2.7.2) \quad \sigma^{p^{s_i}} = \tau \cdot \prod_{j=i+1}^n t_j^{h_j}, \quad \text{with } h_j \in [0, p^{s_i} - 1].$$

It follows from the definition of the inverse Cartier operator, that an $R_i^{p^{s_i}}$ -basis of $\sigma^{p^{s_i}} \cdot B_{s_i} \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$ as submodule of $\tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$ is given by

$$(2.7.3) \quad \sigma^{p^{s_i}} \cdot t_{k_1}^{p^{\mu}-1} \cdots t_{k_{\rho_1}}^{p^{\mu}-1} \cdot dt_{k_1} \cdots dt_{k_{\rho_1}} \cdot d\log t_{l_1} \cdots d\log t_{l_{\rho_2}},$$

where k_ν, l_ν are as above, $\mu \in [0, s_i - 1]$, and $\rho_1 \geq 1, \rho_2 \geq 0$ with $\rho_1 + \rho_2 = r - 1$. By (2.7.2) and direct inspection of the bases above, we find that the inclusion

$$\sigma^{p^{s_i}} \cdot B_{s_i} \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n)) \hookrightarrow \tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$$

is a split-injection of $R_i^{p^{s_i}}$ -modules. Hence $Q_D^{i,r-1}$ is a locally free \mathcal{O}_{D_i} -module. \square

Lemma 2.8. *Let (R, \mathfrak{m}) be a regular local ring and $t_1, \dots, t_n \in \mathfrak{m}$ be part of a regular sequence of local parameters ($n \geq 1$). Set $R_n := R/(t_n)$ and denote by \bar{t}_i the image of t_i in R_n . Then the following sequence is exact*

$$0 \rightarrow 1 + t_n R \rightarrow (R[\frac{1}{t_1 \cdots t_{n-1}}])^\times \rightarrow (R_n[\frac{1}{\bar{t}_1 \cdots \bar{t}_{n-1}}])^\times \rightarrow 0.$$

Furthermore, $1 + t_n R$ is multiplicatively generated by elements $1 + t_n u$, with $u \in R^\times$.

Proof. By the Auslander-Buchsbaum theorem, R is factorial. Since $R/(t_i)$ and $R/(t_n, t_i)$ are again local regular rings (in particular they are integral), we see that

t_1, \dots, t_n are prime elements in R and $\bar{t}_1, \dots, \bar{t}_{n-1}$ are prime elements in R_n . It follows that every unit in $v \in R[\frac{1}{t_1 \cdots t_{n-1}}]$ can be uniquely written in the form

$$v = ut_1^{r_1} \cdots t_{n-1}^{r_{n-1}}, \quad \text{with } u \in R^\times, r_1, \dots, r_{n-1} \in \mathbb{Z},$$

and similar with $R_n[\frac{1}{\bar{t}_1 \cdots \bar{t}_{n-1}}]$. Hence $v \equiv 1 \pmod{t_n}$ implies that $r_1 = \dots = r_{n-1} = 0$ and $u \in 1 + t_n R$. This implies the exactness of the sequence in the statement. For the last part observe that if $a \in \mathfrak{m}$, then $1 + t_n a = (1 + t_n u) \cdot (1 + t_n v)$ with $u = a - 1$, $v = 1/(1 + t_n u) \in R^\times$. \square

Proposition 2.9. *Let the notations and assumptions be as in 2.6. Then there is an isomorphism of sheaves of abelian groups*

$$\theta : Q_D^{i, r-1} \xrightarrow{\simeq} \frac{V_r^i}{V_r^{i-1}}$$

given on local sections by

$$\theta(a \operatorname{dlog} b_1 \cdots \operatorname{dlog} b_{r-1}) = \{1 + \tilde{a}, \tilde{b}_1, \dots, \tilde{b}_{r-1}\},$$

where $\tilde{a} \in \mathcal{O}_X(-D - E_i)$, $\tilde{b}_j \in \mathcal{O}_{X \setminus E_i}^\times$ are local lifts of a and b_j , respectively.

Proof. Note that both sheaves in the statement have support in D_i . Therefore it suffices to show the statement in every point $x \in D_i$. Up to replacing $\{D_j\}_j$ by an ordered subset we may assume that all components of D meet in x . Thus we can assume $X = \operatorname{Spec} R$, where R is a local ring with maximal ideal \mathfrak{m} , which is smooth and essentially of finite type over k . Let $t_j \in \mathfrak{m}$ be a local equation of D_j , $j = 1, \dots, n$. Set $R_i := R/(t_i)$.

We first show that a well-defined map θ as in the statement exists. Set $M^{\text{gp}} := R_i[\frac{1}{t_{i+1} \cdots t_n}]^\times$ and denote by τ the equation (2.7.1) for $D + E_i$. There is a well-defined map

$$(2.9.1) \quad \tau R_i \oplus \bigoplus_{j=1}^{r-1} M^{\text{gp}} \rightarrow \frac{V_r^i}{V_r^{i-1}}$$

given by

$$(a, (b_1, \dots, b_{r-1})) \mapsto \{1 + \tilde{a}, \tilde{b}_1, \dots, \tilde{b}_{r-1}\},$$

where $\tilde{a} \in \tau R$, $\tilde{b}_j \in (R[\frac{1}{t_{i+1} \cdots t_n}])^\times$ are lifts of a and b_j , respectively. We have to check the independence of the choices: Choosing a different lift of a results in multiplying $1 + \tilde{a}$ by an element $(1 + t_i \tau_i c)$, for $c \in R$. The independence of the choice of \tilde{a} follows, since $\{1 + t_i \tau_i c, \tilde{b}\} \in V_r^{i-1}$. Two different lifts of b_1 differ by Lemma 2.8 by a product of element of the form $(1 + t_i e)$ with $e \in R^\times$. Let $c \in R$ and \tilde{b}_j as above, we compute

$$\begin{aligned} \{1 + \tau c, 1 + t_i e, \tilde{b}\} &= -\{1 + et_i, 1 + c\tau, \tilde{b}\} \\ &= -\left\{1 + \frac{ec}{1 + et_i} t_i \tau, -et_i(1 + c\tau), \tilde{b}\right\}, \quad \text{by 2.4(1),} \\ &\in V_r^{i-1}. \end{aligned}$$

This shows the independence of the choice of the lift of \tilde{b}_1 and similar also for \tilde{b}_j . Clearly (2.9.1) is \mathbb{Z} -linear in the b_j ; the linearity in a follows from

$$(1 + \tilde{a}_1)(1 + \tilde{a}_2) = (1 + \tilde{a}_1 + \tilde{a}_2)(1 + t_i c),$$

for $\tilde{a}_i, c \in \tau R$. Furthermore, if $p \neq 2$, then

$$\{1 + \tau c, \tilde{b}\} \equiv 2 \cdot \{1 + \frac{1}{2}\tau c, \tilde{b}\} \pmod{V_r^{i-1}}.$$

Thus the formula $\{b, b\} = \{b, -1\}$ in $K_2^M(k(X))$ implies that (2.9.1) is alternating in the b 's (also for $p = 2$). Altogether we see that (2.9.1) induces a surjective map

$$(2.9.2) \quad \tau R_i \otimes_{\mathbb{Z}} \bigwedge_{j=1}^{r-1} M^{\text{gp}} \twoheadrightarrow \frac{V_r^i}{V_r^{i-1}}.$$

We claim that this map factors via $\tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$. By [Kat89, (1.7)] we have to show that (2.9.2) maps the following elements to zero

$$(2.9.3) \quad \tau a \otimes a \wedge b_1 \wedge \cdots \wedge b_{r-2} - \sum_{\nu} \tau u_{\nu} \otimes u_{\nu} \wedge b_1 \wedge \cdots \wedge b_{r-2},$$

where $a \in M = M^{\text{gp}} \cap R_i$ and $u_{\nu} \in R_i^{\times}$ such that $a = \sum_{\nu} u_{\nu}$, and $b_j \in M^{\text{gp}}$. Choose lifts $\tilde{u}_{\nu} \in R^{\times}$ of u_{ν} and set $\tilde{a} = \sum_{\nu} \tilde{u}_{\nu}$. Let $\tilde{b} = (\tilde{b}_1, \dots, \tilde{b}_{r-2})$ with $\tilde{b}_j \in R[\frac{1}{t_{i+1} \cdots t_n}]^{\times}$ lifts of b_j . Write

$$(1 + \tau \sum_{\nu} \tilde{u}_{\nu}) = \prod_{\nu} (1 + \tau \tilde{u}_{\nu}) \cdot (1 + \tau t_i e),$$

for some $e \in R$. Assume first $e \in R^{\times}$ is a unit. Then in $K_r^M(k(X))$

$$\begin{aligned} \{1 + \tau \tilde{a}, \tilde{a}, \tilde{b}\} &= -\{1 + \tau \tilde{a}, -\tau, \tilde{b}\} \\ &= -\sum_{\nu} \{1 + \tau \tilde{u}_{\nu}, -\tau, \tilde{b}\} - \{1 + \tau t_i e, -\tau, \tilde{b}\} \\ &= \sum_{\nu} \{1 + \tau \tilde{u}_{\nu}, \tilde{u}_{\nu}, \tilde{b}\} + \underbrace{\{1 + \tau t_i e, t_i e, \tilde{b}\}}_{\in V_r^{i-1}}. \end{aligned}$$

Since the starting term and the final term are in V_r^i , the equality of these two terms holds in V_r^i . If e is not a unit we can write

$$(1 + \tau t_i e) = (1 + \tau t_i (e - 1)) \cdot (1 + \tau t_i \frac{1}{1 + \tau t_i (e - 1)})$$

and argue similarly. Hence (2.9.2) sends the relation (2.9.3) to zero and induces a surjection

$$(2.9.4) \quad \tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n)) \twoheadrightarrow \frac{V_r^i}{V_r^{i-1}}.$$

Assume $s_i \geq 1$. Let $\sigma^{p^{s_i}}$ be the equation (2.7.2) for $p^{s_i} D(i)$. We want to show that (2.9.4) maps $\sigma^{p^{s_i}} \cdot B_{s_i} \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))$ to zero. The latter module is generated as abelian group by the elements (cf. [Ill79, 0, Proposition 2.2.8])

$$(2.9.5) \quad (\sigma^{p^{s_i-\nu}} u)^{p^{\nu}} \text{dlog } u \text{dlog } v_1 \cdots \text{dlog } v_{r-2}, \quad u \in R_i^{\times}, v_j \in M^{\text{gp}}, 0 \leq \nu \leq s_i - 1.$$

We compute in $K_r^M(k(X))$ (cf. [BK86, Lemma (4.5)])

$$\begin{aligned} \{1 + (\sigma^{p^{s_i-\nu}} \tilde{u})^{p^{\nu}}, \tilde{u}, \tilde{v}\} &= p^{\nu} \{1 + \sigma^{p^{s_i-\nu}} \tilde{u}, \tilde{u}, \tilde{v}\} \\ &= -p^{\nu} \{1 + \sigma^{p^{s_i-\nu}} \tilde{u}, -\sigma^{p^{s_i-\nu}}, \tilde{v}\} \\ &= -\{1 + \sigma^{p^{2s_i-\nu}} \tilde{u}^{p^{s_i}}, -\sigma, \tilde{v}\} \\ &= \{1 + \sigma^{p^{2s_i-\nu}-1} w_1, w_2, \tilde{v}\}, \quad \text{with } w_j \in R^{\times}, \end{aligned}$$

where the last equality holds by Lemma 2.4(2). Since $p^{2s_i-\nu}-1 \geq 2$, for all $\nu \leq s_i-1$, the last term lies in V_r^{i-1} . Thus (2.9.4) maps the element (2.9.5) to zero. We obtain a surjective morphism θ as in the statement. Set $S_i := R_i[\frac{1}{t_1 \cdots t_i \cdots t_n}]$ (we omit t_i in the denominator). Consider the following diagram

$$\begin{array}{ccc} Q_D^{i,r-1}(R_i) = \frac{\tau \cdot \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))}{\sigma^{p^{s_i}} \cdot B_{s_i} \Omega_{R_i}^{r-1}(\log(t_{i+1} \cdots t_n))} & \xrightarrow{\frac{1}{t_i^{m_i}} \cdot \text{restr.}} & \frac{\Omega_{S_i}^{r-1}}{B_{s_i} \Omega_{S_i}^{r-1}} \\ \downarrow \theta & & \simeq \downarrow (2.3.1) \\ \frac{V_r^i(R_i)}{V_r^{i-1}(R_i)} & \xrightarrow{\text{restriction}} & \frac{V_{r,m_i}(S_i)}{U_{r,m_i+1}(S_i)}. \end{array}$$

The diagram is commutative and the vertical arrow on the right hand side is an isomorphism by Lemma 2.3. We can write R as a directed limit of rings A which are smooth over \mathbf{F}_p and on which $t_1 \cdots t_n$ defines a simple normal crossing divisor. For R replaced by such an A the top horizontal map is injective by Lemma 2.7; since directed limits are exact this holds for R as well. Hence θ is injective, which yields the statement. \square

The following corollary is not used in this paper but we include it here for use in [RS21b].

Corollary 2.10. *Let $X \in \mathbf{Sm}$ with $\dim X = d$ and let D be an effective Cartier divisor with simple normal crossing support. Set $\mathcal{G} := V_{r,X|D}/V_{r,X|D+D_{\text{red}}}$, $r \geq 1$, which we view as a sheaf on $Y = D_{\text{red}}$. Then the natural morphism $H^{d-1}(Y_{\text{Zar}}, \mathcal{G}) \rightarrow H^{d-1}(Y_{\text{Nis}}, \mathcal{G})$ is surjective.*

Proof. With the notation from 2.1 we set

$$\mathcal{F} := \frac{U_{r,X|D+D_{\text{red}}}}{U_{r,X|D+2D_{\text{red}}}} \quad \text{and} \quad \mathcal{H} := \frac{V_{r,X|D}}{U_{r,X|D+D_{\text{red}}}}.$$

We have an exact sequence of Nisnevich sheaves

$$(2.10.1) \quad \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0.$$

By Grothendieck-Nisnevich vanishing we obtain exact sequences for $\tau \in \{\text{Zar}, \text{Nis}\}$

$$H^{d-1}(Y_\tau, \mathcal{F}) \rightarrow H^{d-1}(Y_\tau, \mathcal{G}) \rightarrow H^{d-1}(Y_\tau, \mathcal{H}) \rightarrow 0.$$

By 2.5 and Proposition 2.9 the quotient \mathcal{H} admits a finite decreasing filtration whose successive quotients are coherent \mathcal{O}_Y -modules. Hence the natural morphism $H^{d-1}(Y_{\text{Zar}}, \mathcal{H}) \rightarrow H^{d-1}(Y_{\text{Nis}}, \mathcal{H})$ is an isomorphism. By [RS18, Proposition 2.15, Theorem 2.19] (cf. also 2.13 below) the same holds for \mathcal{F} . This yields the statement. \square

Theorem 2.11. *Let X be a smooth k -scheme of pure dimension d and D an effective divisor on X whose support has simple normal crossings. Let E be a vector bundle of dimension $N+1$ on X and denote by $\pi : \mathbf{P} := \text{Proj}(\text{Sym} E^\vee) \rightarrow X$ the projection from the structure map. Let*

$$\eta^t := c_1(\mathcal{O}_{\mathbf{P}}(1))^t \in H_{\text{Nis}}^t(\mathbf{P}, K_{t,\mathbf{P}}^M) = \text{Hom}_{D(\mathbf{P}_{\text{Nis}})}(\mathbb{Z}, K_{t,\mathbf{P}}^M[t]), \quad t = 0 \dots, N.$$

The map in $D(X_{\text{Nis}})$, the derived category of abelian Nisnevich sheaves on X ,

$$(2.11.1) \quad - \cup \eta^t : \bigoplus_{t=0}^N V_{r-t,X|D}[-t] \xrightarrow{\simeq} R\pi_* V_{r,\mathbf{P}|\pi^*D}, \quad r \geq 0,$$

is an isomorphism. If D_{red} is smooth the same is true for the Zariski site.

Proof. Write $D = \sum_{j=1}^n m_j D_j$ as in 2.5. It suffices to show that $\cup \eta^t$ induces an isomorphism $V_{r-t, X|D} \xrightarrow{\cong} R^t \pi_* V_{r, \mathbf{P}|\pi^* D}$, for all $0 \leq t \leq N$, and $R^t \pi_* V_{r, \mathbf{P}|\pi^* D} = 0$, for all $t > N$. This is a local question and we may therefore assume $\mathbf{P} = \mathbf{P}_X^N$. The corresponding statement for $U_{r, X|D+D_{\text{red}}}$ holds by [RS18, Theorem 2.28] (see also 2.14) below. By Proposition 2.9 we are reduced to show:

- (1) $\cup \text{dlog}(\eta^t) : Q_D^{i, r-t-1} \rightarrow R^t \pi_* Q_{\pi^* D}^{i, r-1}$ is an isomorphism, for all $0 \leq t \leq N$ and $1 \leq i \leq n$;
- (2) $R^t \pi_* Q_{\pi^* D}^{i, r-1} = 0$, for all $t > N$ and all $1 \leq i \leq n$,

where $Q_D^{i, r-1}$ is defined in 2.6. By the definition of $Q_D^{i, r-1}$ and the projection formula, this follows from the corresponding statements for $\Omega_{D_i}^{r-1}(\log E_i)$ and - if $p > 0$ - for $B_s \Omega_{D_i}^{r-1}(\log E_i)$ (where $E_i = D_{i+1} + \dots + D_n$). These are well-known, cf. (2.14.2) and the proof of (2.14.4) (in the case $\text{char}(k) > 0$) below. \square

Definition 2.12. Let the notations and assumptions be as in Theorem 2.11. We define

$$\text{tr}_\pi : R\pi_* V_{N+r, \mathbf{P}|\pi^* D}[N] \rightarrow V_{r, X|D}$$

to be the composition of the natural map $R\pi_* V_{N+r, \mathbf{P}|\pi^* D}[N] \rightarrow R^N \pi_* V_{N+r, \mathbf{P}|\pi^* D}$ followed by the inverse of the isomorphism $-\cup \eta^N : V_{r, X|D} \xrightarrow{\cong} R^N \pi_* V_{N+r, \mathbf{P}|\pi^* D}$.

2.1. Small erratum to [RS18]. We use the opportunity to slightly correct an assumption and a proof from [RS18], see 2.13 and 2.14 below. Please note that these corrections do not have any effect on the rest of *loc. cit.*, in particular not on any of the main results.

2.13. In [RS18, 2.1.4] we consider the following situation: Let X be a smooth and separated scheme over k . Let $\sum_{\lambda \in \Lambda} D_\lambda$ be a simple normal crossing divisor (with smooth components D_λ). For $\mathbf{m} = (m_\lambda)_{\lambda \in \Lambda} \in \mathbb{N}^{|\Lambda|}$ we define $D_{\mathbf{m}} = \sum_{\lambda} m_\lambda D_\lambda$ and set for $\nu \in \Lambda$

$$\text{gr}_{D_{\mathbf{m}}}^\nu K_{r, X}^M := U_{r, X|D_{\mathbf{m}}} / U_{r, X|D_{\mathbf{m}+\delta_\nu}},$$

where $\delta_\nu = (0, \dots, 0, 1, 0, \dots, 0)$ with 1 in the ν 's place. (In *loc. cit.* $U_{r, X|D}$ is denoted by $\mathcal{K}_{r, X|D}^M$.) In the Propositions 2.14, 2.15, Theorem 2.19, and Corollary 2.20 of [RS18] we give formulas for $\text{gr}_{D_{\mathbf{m}}}^\nu K_{r, X}^M$. In *loc. cit.* we allow also some of the m_λ to be zero and in this case the formulas are not correct as written. However, if $m_\lambda \geq 1$, for all $\lambda \in \Lambda$, these formulas are correct and this is the only case used in the rest of [RS18]. (The formulas are used to understand the difference between $U_{r, X|D}$ and $U_{r, X|D_{\text{red}}}$; the latter group can then be studied using Corollary 2.10, which is correct as written, i.e., the m_λ may be zero.)

2.14. Let the notation and assumptions be as in [RS18, Lemma 2.25]. We correct the proof of the isomorphisms

(2.14.1)

$$-\cup \text{dlog}(c_1(\mathcal{O}(1))^i) : \omega_{X|D, \mathbf{m}, \nu}^{q-i} / B_{X|D, r, \mathbf{m}, \nu}^{q-i} \xrightarrow{\cong} R^i \pi_{D, \nu*} (\omega_{P_X|P_D, \mathbf{m}, \nu}^q / B_{P_X|P_D, r, \mathbf{m}, \nu}^q),$$

for $i \geq 0$. To this end replace on page 1013 everything after “Now we prove the statement for $\omega_{\mathbf{m}, \nu}^q$ ” in line 8 until “... from Lemma 2.26 below.” in line 25 by the following:

We have a well-known isomorphism

$$(2.14.2) \quad -\cup \text{dlog}(c_1(\mathcal{O}(1))^i) : \Omega_X^{q-i}(\log D) \xrightarrow{\cong} R^i \pi_{X*} \Omega_{P_X}^q(\log P_D).$$

(Without log-poles this can, e.g., be deduced from [SGA7II, Exp XI, Thm 1.1]; the case with log-poles then follows by considering the degree-wise weight filtration and using [De71, (3.1.5.2)].) Using projection formula and base-change (2.14.2) yields an isomorphism

$$(2.14.3) \quad - \cup \text{dlog}(c_1(\mathcal{O}(1))) : \omega_{X|D, \mathfrak{m}, \nu}^{q-i} \xrightarrow{\cong} R^i \pi_{X*} \omega_{P_X|P_{D, \mathfrak{m}, \nu}}^q, \quad i \geq 0.$$

It remains to show

$$(2.14.4) \quad - \cup \text{dlog}(c_1(\mathcal{O}(1))) : B_{X|D, r, \mathfrak{m}, \nu}^{q-i} \xrightarrow{\cong} R^i \pi_{X*} B_{P_X|P_{D, r, \mathfrak{m}, \nu}}^q, \quad i \geq 0.$$

First case: $\text{char}(k) = 0$. In this case the statement follows by descending induction on q from the exact sequence

$$(2.14.5) \quad 0 \rightarrow Z_{X|D, \mathfrak{m}, \nu}^q \rightarrow \omega_{X|D, \mathfrak{m}, \nu}^q \rightarrow B_{X|D, \mathfrak{m}, \nu}^{q+1} \rightarrow 0,$$

the equality $Z_{X|D, \mathfrak{m}, \nu}^q = B_{X|D, \mathfrak{m}, \nu}^q$ (see [BS19, Lemma 6.2]), and $Z_{X|D, \mathfrak{m}, \nu}^N = \omega_{X|D, \mathfrak{m}, \nu}^N$. (Lemma 2.26 is not needed.)

Second case: $\text{char}(k) = p$. By a limit argument we can reduce to the case that k is finitely generated over \mathbf{F}_p , so that $\Omega_X^q(\log D) = 0$, for q large enough. Thus for $r = 1$ the statement follows by descending induction on q from the exact sequence (2.14.5) together with the exact sequence

$$0 \rightarrow B_{X|D, \mathfrak{m}, \nu}^q \rightarrow Z_{X|D, \mathfrak{m}, \nu}^q \rightarrow \omega_{\mathfrak{m}', \nu}^q \rightarrow 0,$$

coming from the Cartier isomorphism [RS18, Theorem 2.16]. For higher r it follows from the inductive definition of $B_{X|D, r, \mathfrak{m}, \nu}^q$ in [RS18, 2.4.4].

2.2. Comparison of different relative Milnor K -sheaves. We will have to use results from [KS86] in which a slightly different version of the relative Milnor K -theory is used. Here we observe that these different version give rise to the same top degree Nisnevich cohomology, so that the difference won't play a role for us.

2.15. Let X be a k -scheme and $D \subset X$ a nowhere dense closed subscheme. Denote by

$$K_{r, X}^{M, \text{naive}} = (\mathcal{O}_X^\times)^{\otimes_{\mathbb{Z}} n} / J,$$

the naive Milnor K -sheaf on X_{Nis} , where $J \subset (\mathcal{O}_X^\times)^{\otimes_{\mathbb{Z}} r}$ denotes the abelian subsheaf locally generated by elements of the form $a_1 \otimes \dots \otimes a_r$, all $a_i \in \mathcal{O}_X^\times$, and $a_i + a_j = 1$, for some $i \neq j$. By [Ker10, Proposition 10 and Theorem 13] there is a surjective map

$$(2.15.1) \quad K_{r, X}^{M, \text{naive}} \twoheadrightarrow K_{r, X}^M$$

and it is an isomorphism at the stalks x with infinite residue field $k(x)$. In [KS86, (1.3)] the relative Milnor K -sheaf of (X, D) is defined by the formula

$$(2.15.2) \quad K_r^M(\mathcal{O}_X, I) := \text{Ker}(K_{r, X}^{M, \text{naive}} \rightarrow i_* K_{r, D}^{M, \text{naive}}),$$

where I is the ideal sheaf of D and $i : D \hookrightarrow X$ denotes the closed immersion.

Lemma 2.16. *Let the notations and assumptions be as in 2.15. Assume X is noetherian, reduced, and of pure dimension $d < \infty$. Then (2.15.1) induces a surjection*

$$(2.16.1) \quad K_r^M(\mathcal{O}_X, I) \twoheadrightarrow V_{r, X|D}.$$

where the first map is surjective and the second map is injective. Furthermore, the induced map on top-degree Nisnevich cohomology is an isomorphism

$$(2.16.2) \quad H^d(X_{\text{Nis}}, K_r^M(\mathcal{O}_X, I)) \xrightarrow{\cong} H^d(X_{\text{Nis}}, V_{r, X|D}).$$

Proof. The surjection in (2.16.1) holds by [KS86, Lemma 1.3.1]. By [Ker10, Proposition 10(4)] the map (2.15.1) is an isomorphism on fields. Hence (2.16.1) has kernel supported in codimension ≥ 1 . The isomorphism in (2.16.2) follows therefore from Grothendieck-Nisnevich vanishing. \square

3. THE CYLCE CLASS MAP

In this section k denotes a field. For a scheme Z we denote by $Z_{(i)}$ (resp. $Z^{(i)}$) the set of points of Z whose closure have (co)dimension i .

3.1. Let X be a finite type k -scheme and D an effective Cartier divisor on X such that $U = X \setminus |D|$ is smooth over k . For $C \subset X$ an integral curve not contained in the support of D and with normalization $\nu : \tilde{C} \rightarrow C$, we set

$$(3.1.1) \quad G(C, D) := \bigcap_{x \in \tilde{C} \cap \nu^{-1}(D)} \text{Ker}(\mathcal{O}_{\tilde{C}, x}^\times \rightarrow \mathcal{O}_{\tilde{C} \times_C D, x}^\times) \subset k(C)^\times.$$

We define a map

$$\partial_C : G(C, D) \xrightarrow{\text{div}_{\tilde{C}}} Z_0(\tilde{C}) \xrightarrow{\nu_*} Z_0(U),$$

and put

$$\text{CH}_0(X|D) = \text{Coker} \left(\partial = \sum_C \partial_C : \bigoplus_C G(C, D) \rightarrow Z_0(U) \right),$$

where the sum is over all $C \subset X$ as above and $Z_0(Y)$ denotes the group of zero-cycles on Y .

For a closed point $x \in U$ the Gersten resolution ([Ker10, Proposition 10(8)]) yields an isomorphism

$$(3.1.2) \quad \theta_x : \mathbb{Z} \xrightarrow{\cong} H_x^d(U_{\text{Nis}}, K_{d,U}^M) \cong H_x^d(X_{\text{Nis}}, V_{d,X|D}),$$

where $d = \dim X$ and we use the notation from 2.1. By [KS86, Theorem 2.5] and Lemma 2.16 we obtain a surjective map

$$(3.1.3) \quad \theta = \sum_x \theta_x : Z_0(U) = \bigoplus_{x \in U_{(0)}} \mathbb{Z} \twoheadrightarrow H^d(X_{\text{Nis}}, V_{d,X|D}).$$

Similarly, we can define a map $Z_0(U) \rightarrow H^d(X_{\text{Nis}}, U_{d,X|D})$. As a special case of [RS18, Proposition 3.3] this map factors via $\text{CH}_0(X|D)$, if X is smooth and D has simple normal crossing support. We now prove the following refinement of this statement with $U_{d,X|D}$ replaced by $V_{d,X|D}$. In view of (2.16.2), Theorem 1.1 follows from this.

Theorem 3.2. *Let the notation and assumptions be as in 3.1 above. We assume additionally that X is smooth of pure dimension d and that the support of D is a simple normal crossing divisor. Then (3.1.3) factors to give a surjective map*

$$\text{CH}_0(X|D) \twoheadrightarrow H^d(X_{\text{Nis}}, V_{d,X|D}).$$

Remark 3.3. The above factorization of (3.1.3) via $\text{CH}_0(X|D)$ is proven for quasi-projective surfaces in [BKS21] and for smooth projective schemes over a finite field in [GK21], see Theorem 1.3. In both cases the divisor D is allowed to be an arbitrary effective Cartier divisor.

Proof. The proof is similar to [RS18, Proposition 3.3], though we will need the projective bundle formula for $V_{r,X|D}$ proved in the previous section and we take a short cut around the Cousin resolution used in *loc. cit.*

We have the spectral sequence

$$E_1^{i,j} = \bigoplus_{x \in X^{(i)}} H_x^{i+j}(X_{\text{Nis}}, V_{d,X|D}) \implies H^{i+j}(X_{\text{Nis}}, V_{d,X|D}).$$

Grothendieck vanishing ([Nis89, Corollary 1.3.3]) yields

$$\begin{aligned} H^d(X_{\text{Nis}}, V_{d,X|D}) &= E_2^{d,0} \\ &= \text{Coker} \left(\bigoplus_{x \in X_{(1)}} H_x^{d-1}(X_{\text{Nis}}, V_{d,X|D}) \xrightarrow{\partial} \bigoplus_{x \in X_{(0)}} H_x^d(X_{\text{Nis}}, V_{d,X|D}) \right). \end{aligned}$$

Assume $C \subset X$ is a regular curve not contained in D and $f \in k(C)^\times$ with $f \equiv 1 \pmod{D|_C}$. We claim that

$$\text{div}(f) \in \bigoplus_{x \in U_{(0)}} \mathbb{Z} \subset \bigoplus_{x \in X_{(0)}} H_x^d(X_{\text{Nis}}, V_{d,X|D}),$$

lies in the image of

$$(3.3.1) \quad \partial = (\partial_x)_x : k(C)^\times \cong H_c^{d-1}(X_{\text{Nis}}, V_{d,X|D}) \rightarrow \bigoplus_{x \in X_{(0)}} H_x^d(X_{\text{Nis}}, V_{d,X|D}),$$

where c is the generic point of C . We have $\partial(f)|_U = \pm \text{div}_{C \cap U}(f)$ (e.g. [RS18, (3.3.1)]). Thus it remains to show $\partial_x(f) = 0$, for all $x \in D_{(0)}$. To this end it suffices to show that f lies in the image

$$(3.3.2) \quad H_{C_{(x)}}^{d-1}(\text{Spec } \mathcal{O}_{X,x}, V_{d,X|D}) \rightarrow H_c^{d-1}(X, V_{d,X|D}),$$

for $x \in D_{(0)}$, where $C_{(x)} = \text{Spec } \mathcal{O}_{C,x}$. Under the isomorphism

$$k(C)^\times \cong H_c^{d-1}(X, V_{d,X|D}) = H_c^{d-1}(X, K_{d,X}^M)$$

f corresponds by [RS18, Corollary 2.3] (and with the notation from there) to

$$(3.3.3) \quad \begin{bmatrix} \{\tilde{f}, t_1, \dots, t_{d-1}\} \\ t_1, \dots, t_{d-1} \end{bmatrix},$$

where $\tilde{f} \in \mathcal{O}_{X,c}$ is a lift of f and $t_1, \dots, t_{d-1} \in \mathfrak{m}_c$ is a regular system of parameters. In fact since C and X are regular, the closed immersion $C \hookrightarrow X$ is regular and hence we can choose the t_i 's to be a regular sequence in $\mathcal{O}_{X,x}$ generating the ideal sheaf of C at x . Furthermore, since f satisfies the modulus condition we can choose the lift \tilde{f} to lie in $\mathcal{O}_{X|D,x}^\times$. This shows that (3.3.3) lies in the image of (3.3.2). Thus we have shown the above claim that $\text{div}(f)$ lies in the image of (3.3.1), if $C \subset X$ is regular.

Assume $C \subset X$ is an arbitrary integral curve not contained in D . Let $\nu : \tilde{C} \rightarrow C$ be the normalization. We embed \tilde{C} in $P_X = \mathbf{P}_X^n$ over X . Denote by $\pi : P_X \rightarrow X$

the projection and by π_U its base change over U . Consider the following diagram

$$\begin{array}{ccccc} Z_0(\tilde{C} \cap P_U) & \longrightarrow & H_{(P_U)_{(0)}}^{d+n}(P_U, \text{Nis}, K_{d+n, P_U}^M) & \longrightarrow & H^{d+n}(P_{X, \text{Nis}}, V_{d+n, P_X|P_D}) \\ \downarrow \nu_* & & \downarrow \text{tr}_{\pi_U} & & \downarrow \text{tr}_{\pi} \\ Z_0(C \cap U) & \longrightarrow & H_{U_{(0)}}^d(U, \text{Nis}, K_{d, U}^M) & \longrightarrow & H^d(X_{\text{Nis}}, V_{d, X|D}), \end{array}$$

where tr_{π} and tr_{π_U} are defined in 2.12 and we view $U_{(0)}$ and $(P_U)_{(0)}$ as families of closed supports on U and P_U , respectively. The left square commutes by [RS18, Lemma 2.27], the right square by construction, and the composite horizontal maps are equal to (3.1.3). Since \tilde{C} is regular the case considered above yields that $\text{div}_{\tilde{C}}(f)$ is mapped to zero in $H^{d+n}(P_{X, \text{Nis}}, V_{d+n, P_X|P_D})$, for $f \equiv 1 \pmod{D|_{C'}}$. Hence $\text{div}_C(f) = \nu_* \text{div}_{\tilde{C}}(f)$ is mapped to zero in $H^d(X_{\text{Nis}}, V_{d, X|D})$. This completes the proof. \square

Remark 3.4. The assumption that the support of D is a simple normal crossing divisor, was only used to apply the projective bundle formula, Theorem 2.11.

REFERENCES

- [BK86] S. Bloch and K. Kato, *p-adic étale cohomology*, Institut des Hautes Études Scientifiques. Publications Mathématiques, Inst. Hautes Études Sci. Publ. Math, (1986) **63**, 107–152.
- [BKS21] F. Binda, A. Krishna, and S. Saito, *Bloch’s formula for 0-cycles with modulus and higher dimensional class field theory*, to appear in JAG, <https://arxiv.org/abs/2002.01856>.
- [BS19] F. Binda and S. Saito, *Relative cycles with moduli and regulator maps*, J. Inst. Math. Jussieu **18** (2019), no. 6, 1233–1293.
- [De71] P. Deligne, *Théorie de Hodge. II*, Inst. Hautes Études Sci. Publ. Math., Institut des Hautes Études Scientifiques. Publications Mathématiques, **40** (1971), 5–57.
- [GK21] R. Gupta and A. Krishna, *Idele class groups with modulus*, Preprint 2021, <https://arxiv.org/abs/2101.04609>.
- [GK20] R. Gupta and A. Krishna, *K-theory and 0-cycles on schemes*, J. Algebraic Geom., **29** (2020), no. 3, 547–601.
- [Kat89] K. Kato, *Logarithmic structures of Fontaine-Illusie*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988) (1989), 191–224, Johns Hopkins Univ. Press, Baltimore, MD.
- [Ill79] L. Illusie, *Complexe de de Rham-Witt et cohomologie cristalline*, Ann. Sci. École Norm. Sup. (4) **12** (1979), no. 4, 501–661.
- [KS86] K. Kato and S. Saito, *Global class field theory of arithmetic schemes*, Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), Contemp. Math., vol. 55, Amer. Math. Soc., Providence, RI, 1986, pp. 255–331. MR 862639
- [Ker10] M. Kerz, *Milnor K-theory of local rings with finite residue fields*, J. Algebraic Geom. **19** (2010), no. 1, 173–191.
- [KS16] M. Kerz and S. Saito, *Chow group of 0-cycles with modulus and higher-dimensional class field theory*, Duke Math. J. **165** (2016), no. 15, 2811–2897. MR 3557274
- [KS14] M. Kerz and S. Saito, *Lefschetz theorem for abelian fundamental group with modulus*, Algebra Number Theory **8** (2014), no. 3, 689–701.
- [Kri15] A. Krishna, *On 0-cycles with modulus*, Algebra Number Theory **9** (2015), no. 10, 2397–2415.
- [Kri18] A. Krishna, *Torsion in the 0-cycle group with modulus*, Algebra Number Theory, **12** (2018), no. 6, 1431–1469.
- [Nis89] Y. A. Nisnevich, *The completely decomposed topology on schemes and associated descent spectral sequences in algebraic K-theory*, Algebraic K-theory: connections with

- geometry and topology (Lake Louise, AB, 1987), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 279, Kluwer Acad. Publ., Dordrecht, 1989, pp. 241–342.
- [RS21a] K. Rülling and S. Saito, *Ramification theory of reciprocity sheaves, I, Zariski-Nagata Purity*. In preparation.
- [RS21b] K. Rülling and S. Saito, *Ramification theory of reciprocity sheaves, II, Higher local symbols*. In preparation.
- [RS18] K. Rülling and S. Saito, *Higher Chow groups with modulus and relative Milnor K -theory*, Trans. Amer. Math. Soc. **370** (2018), no. 2, 987–1043.
- [SGA7II] *Groupes de monodromie en géométrie algébrique. II*, Lecture Notes in Mathematics, Vol. 340, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 II), Dirigé par P. Deligne et N. Katz, Springer-Verlag, Berlin-New York, (1973).

BERGISCHE UNIVERSITÄT WUPPERTAL, GAUSSSTR. 20, D-42119 WUPPERTAL, GERMANY
Email address: ruelling@uni-wuppertal.de

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA,
TOKYO 153-8941, JAPAN
Email address: sshuji@msb.biglobe.ne.jp